

SHORT NOTE

Predation on *Amerotyphlops brongersmianus* (Squamata, Typhlopidae) by *Bothrops moojeni* (Squamata, Viperidae) in west Brazil

Depredación de *Amerotyphlops brongersmianus* (Squamata, Typhlopidae) por *Bothrops moojeni* (Squamata, Viperidae) en el oeste de Brasil

José Gonçalves De Oliveira-Jr.¹ | Karoline Ceron^{1*} | Sarah Mângia¹ | Isabela C. O Da Silva² | José Luiz Massao Moreira Sugai¹ | Diego J. Santana¹

- Received: 17/oct/2018
- Accepted: 13/aug/2019
- Published online: 29/oct/2019

Citation: Oliveira-Jr JG, Ceron K, Mângia S, Silva ICO, Sugai JLMM, Santana DJ. 2020 Predation on *Amerotyphlops brongersmianus* (Squamata, Typhlopidae) by *Bothrops moojeni* (Squamata, Viperidae) in West Brazil. *Caldasia* 42(1):150-152. doi: <https://dx.doi.org/10.15446/caldasia.v42n1.74998>.

ABSTRACT

Ophyophagy is a common feeding habit in snakes; however, there are few records of this behavior for the genus *Bothrops*. Here, we report the first case of predation of *Bothrops moojeni* upon *Amerotyphlops brongersmianus*. Our record reinforces the known generalist habit of *Bothrops moojeni* and indicates an opportunistic diet.

Keywords. Diet, ophyophagy, pitvipers, snakes, wormsnakes

RESUMEN

Ofofagia es un hábito alimenticio común en serpientes, sin embargo, hay pocos registros de este comportamiento para el género *Bothrops*. En el presente trabajo reportamos el primer caso de depredación de *Bothrops moojeni* en *Amerotyphlops brongersmianus*. Nuestro registro refuerza el conocimiento del hábito generalista de *Bothrops moojeni* e indica una dieta oportunista de esta especie.

Palabras clave. Dieta, ofiofagia, serpientes, serpientes-gusano, víboras

¹ Universidade Federal de Mato Grosso do Sul, Instituto de Biociências, Cidade Universitária, CEP 79002-970, Campo Grande, Mato Grosso do Sul, Brazil, jose_jr_96@hotmail.com, adenomera@gmail.com*, sarahmangia@yahoo.com.br, jlmassao@gmail.com, santanadiejojose@yahoo.com

² Universidade Federal de Mato Grosso do Sul, Laboratório de Parasitologia Animal, Cidade Universitária, CEP 79070-900, Campo Grande, MS, Brazil, isabela.caroline@hotmail.com

* Corresponding author



All the species of snakes are carnivorous, and most of them are generalists regarding the prey (Martins *et al.* 2002). Ophiophagy, when a species feeds on snakes, is not a rare feeding habit in snakes; such is the case of the mussuranas (*Clelia* spp.) and the king cobra (*Ophiophagus hannah* Cantor, 1836) (Marques *et al.* 2016).

Bothrops moojeni Hoge, 1966 is a nocturnal, large pit viper distributed in central and southeastern Brazil, throughout riparian forests and adjacent open areas in the Cerrado morphoclimatic domain (Campbell and Lamar 2004). Its diet is composed mainly by mammals, frogs, and lizards (Nogueira *et al.* 2003). Nogueira *et al.* (2003) also recorded snakes on the *B. moojeni* diet; however, the authors did not specify the identity of the preyed species. Here, we report the first case of predation of *Bothrops moojeni* upon two individuals of *Amerotyphlops brongersmianus* (Vanzolini, 1976), a fossorial snake widely distributed in South America, inhabiting various habitats (Dixon and Hendricks 1979).

On 1^o February 2018, during a fieldwork at Parque Estadual das Nascentes do Rio Taquari (18°15'87" South,

53°41'767" West, 450 m) in Alcinoópolis municipality, Mato Grosso do Sul state, Midwest Brazil, we captured an adult of *B. moojeni* (ZUFMS-REP 3318, SVL = 750 mm, tail = 170 mm) (Collection permit SISBIO #45889/1 and IMASUL #71/400151/2018). During the analysis of parasitism in the body of *B. moojeni*, we found two individuals of *A. brongersmianus* in the stomach of *B. moojeni* (ZUFMS-REP 3319, SVL = 60.68 mm and ZUFMS-REP 3320, SLV = 120.32 mm) (Fig. 1). All the three snakes are housed at Coleção Zoológica da Universidade Federal de Mato Grosso do Sul.

There are few records of ophiophagy for the genus *Bothrops* (e.g. *B. leucurus* – Fagundes *et al.* 2009; *B. atrox* - Oliveira and Martins 1996; *B. jararacussu* - De Roodt and Lanari 2015; *B. moojeni* – Nogueira *et al.* 2003). Previous studies indicate that the genus is composed by generalist species, feeding on small rodents, amphibians, lizards, and fish (*B. jararaca* - Hartmann *et al.* 2009, *B. atrox* - Bernarde and Abe 2010).

Bothrops moojeni also shows an ontogenetic diet shift, varying from ectothermic to endothermic prey throughout



Figure 1. Two individuals of *Amerotyphlops brongersmianus* (SVL = 60.68 mm, ZUFMS-REP 3319 and SLV = 120.32 mm, ZUFMS-REP 3320) preyed by an adult of *Bothrops moojeni* (SVL = 750 mm, ZUFMS-REP 3318).

its life (Nogueira et al. 2003), with adults preying more frequently on mammals (França et al. 2008). This is the first record of *A. brongermianus* as prey of *B. moojeni*. Besides our observation, this fossorial snake was already reported as prey of *Bothrops leucurus* Wagler, 1824 and *Micrurus ibiboboca* (Merrem, 1820) (Baptista et al. 2016, Marques et al. 2016). Our record reinforces the known generalist habit of *B. moojeni* and indicates an opportunistic diet, with the snake feeding on preys that share and inhabits the same habitat.

LITERATURE CITED

- Baptista GM, Gonzalez RC, Marcial De Castro T, Silva-Soares T. 2016. *Bothrops leucurus* (White-tailed Lancehead). DIET. Herpetol. Rev. 47(3):474–475.
- Bernarde OS, Abe AS. 2010. Hábitos alimentares de serpentes em Espigão do Oeste, Rondônia, Brasil. *Biota Neotrop.* 10(1):167–173. doi: <https://doi.org/10.1590/S1676-06032010000100017>.
- Campbell JA, Lamar WW. 2004. *The Venomous Reptiles of the Western Hemisphere*. New York: Cornell University Press.
- De Roodt AR, Lanari LC. 2015. Notes of cases of aggression and ophiophagy by *Bothrops jararacussu* (Lacerda 1884) in captivity. *Acta toxicol. Argent.* 23(3):134–141.
- Dixon JR, Hendricks FS. 1979. The wormsnakes (Family Typhlopidae) of the neotropic, exclusive of the Antilles. *Zool. Verh.* 173:1–39.
- Fagundes V, Baião LA, Vianna LA, Alvarenga CS, Machado MX, Lopes, SR. 2009. *Bothrops leucurus* (Serpentes, Viperidae) preying on *Micrurus corallinus* (Serpentes, Elapidae) and *Blarinomys breviceps* (Mammalia, Cricetidae). *Bol. Mus. Biol. Mello Leitão.* 25:67–71.
- França FGR, Mesquita DO, Nogueira CC, Araújo AFB. 2008. Phylogeny and ecology determine morphological structure in a snake assemblage in the Central Brazilian Cerrado. *Copeia* 2008(1):23–38. doi: <https://doi.org/10.1643/CH-05-034>.
- Hartmann PA, Hartmann MT, Martins M. 2009. Ecologia e história natural de uma taxocenose de serpentes no Núcleo Santa Virgínia do Parque Estadual da Serra do Mar, no sudeste do Brasil. *Biota Neotrop.* 9(3):173–184. doi: <https://doi.org/10.1590/S1676-06032009000300018>.
- Marques R, Mebert K, Fonseca É, Rödder D, Solé M, Tinôco M. S. 2016. Composition and natural history notes of the coastal snake assemblage from Northern Bahia, Brazil. *ZooKeys* 611:93–142. doi: <https://doi.org/10.3897/zookeys.611.9529>.
- Martins M, Marques OAV, Sazima I. 2002. Ecological and phylogenetic correlates of feeding habits in Neotropical pit vipers of the genus *Bothrops*. In: Schuett GW, Höggren M, Douglas ME, Greene HW, editors. Chapter 20, *Biology of the Vipers*. Eagle Mountain: Eagle Mountain Publishing; p.307–328.
- Nogueira C, Sawaya RJ, Martins M. 2003. Ecology of the pitviper, *Bothrops moojeni*, in the Brazilian Cerrado. *J. Herpetol.* 37(4):653–659. doi: <https://doi.org/10.1670/120-02A>.
- Oliveira ME, Martins M. 1996. *Bothrops atrox* (Common Lancehead) foraging behavior and ophiophagy. *Herpetol. Rev.* 27(1):22–23.

AUTHOR'S CONTRIBUTION

KC, ICO, JLMMS and DJS collected data; JGOJr and SM led the writing with contributions from all authors, interpretation and discussion of the results.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.