http://doi.org/10.17648/heringeriana.v17i1.918025

Heringeriana 17: e918025. 2023.
http://revistas.jardimbotanico.ibict.br/
Jardim Botânico de Brasília

ISSN 1983-6996 Versão impressa
Heringeriana
ISSN 2359-165X Versão *on-line*

Scientific note

Predation of *Boana faber* (Anura, Hylidae) upon *Dendropsophus minutus* (Anura, Hylidae)

Henrique J. Oliveira[1*], Márcia M. P. Müller[2], Priscila S. Carvalho[2,3] & Diego J. Santana[3]

The Blacksmith Tree Frog *Boana faber* (Wied-Neuwied, 1821) is a large-sized species that inhabits water bodies in forested and open environments in the Atlantic Forest, from northeastern Argentina to eastern Brazil (Faivovich *et al.* 2005; Haddad *et al.* 2013). The vocalization of this species resembles the hammering of a blacksmith and males have a prepollex with a spine resembling a thumb, which is used in combat with other males. The *Boana faber* reproduction involves mud nests constructed by males at the edges of bodies of water (Martins, 1993; Maffei & Ubaid, 2014). Despite some aspects of its biology are well known, like his breeding behavior (Martins, 1993) and movement (Oliveira *et al.* 2016). Observations on its trophic ecology remain scarce, with some information about attempted predation (Rocha-Lima *et al.* 2018) and diet (Table 1). In this contribution we report a predation event by *Boana faber* on November 3rd, 2021, at approximately 8:00 p.m, in Serra do Brigadeiro, municipality of Ervália, state of Minas Gerais, Brazil (-20.86661º S, -42.52393º W, *datum* WGS84; 1,091 m a.s.l [above sea level]). On the ground, on the shores of a small water bodie, we found a male of *B. faber* (ZUFMS-AMP16860) preying upon a *Dendropsophus minutus* (Peters, 1872) (ZUFMS-AMP16859) at the edge of a pond (Figure 1). Both individuals were collected (under collection permit SISBIO #45889) and deposited in the Zoological Collection from Universidade Federal do Mato Grosso do Sul (ZUFMS-AMP).

In addition, aiming to search for records of batracophagy by *B. faber*, we conducted on May 9th, 2022 a literature review using the Google Scholar database, employing the following terms: dieta OR diet AND "*Boana faber*" OR "*Hyla faber*" OR "*Hypsiboas faber*". We found 495 results, of which only four were relevant to this study as they presented batracophagy records by *B. faber*. Additionally, we searched in all editions of the journal Herpetological Review (1967–2022), which traditionally publishes natural history notes, many of which are not yet traceable by Google Scholar. In this journal, we found three records useful to this work, none of which were available on Google Scholar. Therefore, we found a total of nine species preyed upon by *B. faber*, including even a record of cannibalism (Table 1). Once *Dendropsophus minutus*

[1] Programa de Pós-Graduação em Biodiversidade e Conservação da Natureza, Universidade Federal de Juiz de Fora. CEP 36036-900, Juiz de Fora, Minas Gerais, Brazil.
[2] Programa de Pós-graduação em Biodiversidade, Universidade Estadual Paulista "Júlio de Mesquita Filho". CEP 15054-000, São José do Rio Preto, São Paulo, Brazil.
[3] Mapinguari - Laboratório de Sistemática e Biogeografia de Anfíbios e Répteis, Instituto de Biociências, Universidade Federal de Mato Grosso do Sul. CEP 79070-900, Campo Grande, Mato Grosso do Sul, Brazil.
* Corresponding author: henrique.bio22@gmail.com

and *Boana faber* are often found sintopically (e.g., Moura *et al.* 2012; Santana et al., 2010), this interaction might be more frequent along the species distribution. Besides, the Eltonian shortfall refers to a lack of understanding of the ecological interactions between species within an ecosystem, the present record increases our knowledge about the trophic ecology of the smith-frog and its prey.

Table 1. List of amphibians recorded as prey of *Boana faber*.

Preys	**Sources**
Aplastodiscus perviridis (Lutz, 1950)	Solé *et al.* (2004)
Scinax granulatus (Peters, 1871)	Solé *et al.* (2004)
Haddadus binotatus (Spix, 1824)	Leite *et al.* (2008)
Scinax aff. *perereca*	Moura & Feio (2010)
Dendropsophus meridianus (Lutz, 1954)	Figueiredo-de-Andrade *et al.* (2012)
Boana faber	Maffei *et al.* (2014)
Boana semilineata (Spix, 1824)	Braun *et al.* (2017)
Boana pardalis (Spix, 1824)	Braun *et al.* (2017)
Dendropsophus minutus	Present work

Figure 1. Sequence of photos of the *Dendropsophus minutus* being pulled out after it was preyed upon by the *Boana faber*, on November 3rd, 2021 in the Serra do Brigadeiro, municipality of Ervália, Minas Gerais, Brazil.

ACKNOWLEDGMENTS

We are grateful to Dr. Renato N. Feio for allowing access to their property for data collection; to two anonymous referee and editor in chief Dra. Maria R. V. Zanatta for their invaluable feedback and suggestions on earlier drafts of this text. This study is partially funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) – Finance Code 001 (Master's scholarship to HJO and Ph.D. scholarship to MMPM and PSC). DJS thanks CNPq for his research fellowship (CNPq 309420/2020-2).

REFERENCES

Braun, D.V., Lopes, S.R. & Mônico, A.T. (2017) *Hypsiboas faber* (Perereca-martelo; Smith's Frog), Diet. *Herpetological Review* 48: 1–162.

Faivovich, J., Haddad, C.F., Garcia, P.C., Frost, D.R., Campbell, J.A. & Wheeler, W.C. (2005) Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of natural History* 2005: 1–240.

Figueiredo-de-Andrade, C.A., Caram, J. & Carvalho-e-Silva, S.P. (2012) Frog eats frog: Report of three cases from the Atlantic rain forest, southeastern Brazil. *Salamandra* 48: 230–232.

Haddad, C.F., Toledo, L.F., Prado, C.P., Loebmann, D., Gasparini, J.L. & Sazima, I. (2013) *Guia dos anfíbios da Mata Atlântica: diversidade e biologia.* Anolis books, São Paulo, 544 pp.

Hortal, J., Bello, F., Diniz-Filho, J.A.F., Lewinsohn, T.M., Lobo, J.M. & Ladle, R.J. (2015) Seven shortfalls that beset large-scale knowledge of biodiversity. *Annual Review of Ecology, Evolution, and Systematics* 46: 523–549. https://doi.org/10.1146/annurev-ecolsys-112414-054400

Leite, F.S.F., Pezzuti, T.L. & Drummond, L.O. (2008) *Hypsiboas faber* (Smith Frog), Diet. *Herpetological Review* 39: 206–207.

Maffei, F., Nascimento, B.T.M. & Bernarde, P.S. (2014) Cannibalism in the Smith Frog, *Hypsiboas faber* (Wied-Neuwied, 1821), in southern Brazil. *Reptiles & Amphibians* 21: 133–135.

Maffei, F. & Ubaid, F.K. (2014) *Amphibians of Rio Claro Farm, Lençóis Paulista, São Paulo, Brazil.* Editora Canal. 146 pp.

Martins, M. (1993) Observations on the reproductive behaviour of the Smith Frog *Hyla Jaber*. *Herpetological Journal* 3: 31–34.

Moura, M.R. & Feio, R.N. (2010) *Hypsiboas faber* (Wied-Neuwied) predation on *Scinax* aff. *perereca* (Anura: Hylidae) in southeastern Brazil. *Herpetology Notes* 3: 353–354.

Moura, M.R., Motta, A.P., Fernandes, V.D. & Feio, R.N. (2012) Herpetofauna da Serra do Brigadeiro, um remanescente de Mata Atlântica em Minas Gerais, sudeste do Brasil. *Biota Neotropica* 12: 209–235. https://doi.org/10.1590/S1676-06032012000100017

Oliveira, M., Aver, G.F., Moreira, L.F.B., Colombo, P. & Tozetti, A.M. (2016) Daily movement and microhabitat use by the Blacksmith Treefrog *Hypsiboas faber* (Anura: Hylidae) during the breeding season in a subtemperate forest of Southern Brazil. *South American Journal of Herpetology* 11: 89–97. http://dx.doi.org/10.2994/SAJH-D-16-00017.1

Rocha-Lima, A.B.C., Santos, I., Duarte, L.S.C. & da Costa, W.P. (2018) *Erythrolamprus miliaris orinus* (Reptilia, Squamata, Dipsadidae): tentativas de predação de *Boana faber* e *Leptodactylus latrans*. *Boletim do Museu Paraense Emílio Goeldi-Ciências Naturais* 13: 455–460.

Santana, D.J., São Pedro, V.A., Hote, P.S., Roberti, H.M., Sant'Anna, A.C., Figueiredo-de-Andrade, C.A. & Feio, R.N. (2010) Anurans in the region of the High Muriaé River, state of Minas Gerais, Brazil. *Herpetology Notes* 3: 1–10.

Solé, M., Pelz, B. & Kwet, A. (2004) *Hyla faber* (Smith Frog), Diet. *Herpetological Review* 35: 159.

Solé, M., Beckmann, O., Pelz, B., Kwet, A. & Engels, W. (2005) Stomach-flushing for diet analysis in anurans: an improved protocol evaluated in a case study in Araucaria forests, southern Brazil. *Studies on Neotropical Fauna and Environment* 40: 23–28. https://doi.org/10.1080/01650520400025704

Received 14/03/2023
Accepted 31/03/2023
Published 08/05/2022

This is an open-access article distributed under the terms of the Creative Commons Attribution License.