

REVIEW

Open Access

Revisiting the role of anurans in the trophic ecology of Brazilian ecosystems



Camila Fernanda Moser^{1*}, Karoline Ceron², Laura Kauer Schuck³, Diego José Santana⁴, Mirco Solé^{5,6}, Samanta Iop⁷ and Alexandro Marques Tozetti^{8,9}

*Correspondence:
Camila Fernanda Moser
camila-moser@hotmail.com

Full list of author information is
available at the end of the article

Abstract

Amphibians are key players in ecosystem dynamics, and their diets offer critical insights into trophic interactions, community structure, and ecological functioning. However, there is a lack of understanding the role of anurans in the trophic ecosystem dynamics. Here, we gathered available information from the literature to outline current knowledge of the diet of Brazilian post-metamorphic anurans. We retrieved 221 studies spanning from 1971 to June 2025, resulting in dietary data for 233 species, around 19% of Brazilian frog species. Most studies focused on the family Hylidae (84 spp.; 36.1%), followed by Leptodactylidae (55 spp.; 26.6%), Bufonidae (23 spp.; 9.9%), and Microhylidae (15 spp.; 6.4%). Only 12 (1.05%) species classified as threatened to some degree had their diet described. Among Brazilian ecoregions, the Atlantic Forest had the highest number of dietary studies published (110; 49.8%), followed by Amazon (47; 21.3%), Cerrado (32; 14.5%), Caatinga (23; 10.4%), Pampa (10; 4.5%) and Pantanal (8; 3.6%). Despite the increase in diet studies in recent decades, more than 80% of Brazilian anuran species lack dietary data, a trend potentially stemming from the current undervaluation of natural history studies. This scenario is concerning, since anurans are the most threatened vertebrates globally, and information about the natural history of species is fundamental in all phases of research (e.g. development of hypotheses, interpretation, and discussion of data). Our findings underscore a significant information gap in the trophic ecology of Brazilian anurans.

Keywords Ecoregion, Feeding ecology, Food resources, Natural history, Trophic ecology

1 Introduction

Understanding the natural history of species such as diet, behavior, and geographic distribution is essential for experimental design, and for the development and testing of ecological hypotheses [1]. Among them, diet is a core component that provides fundamental information about community dynamics, ecological interactions and ecosystem functioning [2–5]. Dietary analysis can reveal foraging strategies [6, 7] and gain insights into the trophic organization of ecosystems by examining species interactions, including predator-prey dynamics and intra- and interspecific competition [8–10]. Moreover,



combining diet studies with potential prey data from the studied environments can unveil specialized feeding habits and food preferences [3, 6, 7, 11]. When available for sympatric species within the same temporal window, this information enables the inference of trophic niche overlap and resource partitioning strategies [2, 12].

Despite the paramount importance of interactions between species, the Eltonian shortfall persists as a prominent knowledge gap in biodiversity research, particularly in tropical regions [13], which boast unparalleled biodiversity. Brazil, for instance, harbors the world's richest anuran fauna, representing 14.9% of Anura global [14], the lack of comprehensive literature reviews on anuran diets is noticeable. This gap of knowledge presents a paradoxical scenario where a region abounding with biodiversity have few crucial information on species interactions. Hence, elucidating the extent of our knowledge regarding anuran diets is imperative for addressing this deficiency and gaining a deeper understanding of tropical ecosystem dynamics.

Currently, Brazil harbors 1,144 species of anuran amphibians, about 77% of which are endemic to their respective ecoregions (i.e. Amazon, Atlantic Forest, Caatinga, Cerrado, Pampa, and Pantanal) [15, 16]. Furthermore, anuran amphibians are the most endangered group of vertebrates worldwide [17, 18], and specifically in Brazil, they face a great increase in the percentage of threatened species from 3% (35 spp.) in 2004 to 16% (184 spp.) in 2022 [17]. However, for species classified as data-deficient or not assessed, the percentage dropped significantly from 48.7% (557 spp.) to 6.7% (77 spp.) [17]. Despite it, a substantial information gap persists regarding the natural history of these amphibians, resulting in a lack of information for decision-making on the conservation of these species. In 2006, McCallum and McCallum [19] suggested a lack of documentation in natural history in general in Brazilian amphibians. More than a decade later, we see that this gap remains, as documented in the review by Guerra et al. [18], which shows, for example, that almost half of the Brazilian species lack documented vocalizations. Considering that the reproductive behavior of amphibians is one of the most documented across literature [16, 20–22], other areas of their natural history might have even larger gaps.

Beyond taxonomic coverage, it is also essential to consider geographic representativeness in natural history studies. Ecological traits such as diet are strongly influenced by local resource availability, which can vary widely across geographic and environmental gradients [9, 10]. Another important factor to take into consideration is the scientific journals where these studies are published. The identification of journals publishing dietary studies serves as a proxy for evaluating how widely such knowledge is being disseminated across scientific audiences. Thus, identifying spatial gaps in dietary research is key to understanding how anurans interact with their environments across Brazil's heterogeneous landscapes.

Given the importance of understanding anuran diets and the lack of studies on this topic in Brazil, this work aimed to gather and synthesize the information available in literature on the diet of Brazilian anuran amphibians. More specifically, this study aimed to: (i) review the history of dietary research on Brazilian post-metamorphic anurans; (ii) catalog studied species; (iii) analyze publishing journals; (iv) evaluate dietary data for threatened species, and (v) identify the ecoregions with the largest number of studies. Thus, our goal was to pinpoint critical knowledge gaps in this field.

2 Methodology

To ensure a systematic and comprehensive review of the literature on the diet of Brazilian anurans, we followed the PRISMA framework for Systematic Reviews and Meta-Analyses [23]. We searched for studies on the diet of anurans in Brazil in four databases: the Web of Science database (www.webofscience.com), Scientific Electronic Library online (Scielo; www.scielo.org), Google Scholar (www.scholar.google.com) and Scopus (www.scopus.com). For the bibliographic review, we used the following keywords: “Anura” and “diet” or “trophic ecology” or “feeding ecology” or “food habits” or “autoecology” or “trophic overlap” or “trophic composition” and “Brazil”. We also used the same set of words in Portuguese: “Anura” and “dieta” or “ecologia trófica” or “ecologia alimentar” or “hábito alimentar” or “autoecologia” or “sobreposição trófica” or “composição trófica” and “Brasil”. The identified studies were evaluated based on predefined inclusion and exclusion criteria, ensuring the selection of relevant and methodologically sound articles. Studies with taxa identified only at the genus level or those lacking clear species identification (e.g. aff., cf., and gr.) were excluded from consideration, as well as studies on tadpole diet and gray literature (i.e. abstracts presented at conferences, theses, dissertations and other works not published in scientific journals). Furthermore, notes of a single predation or observation of predation were not considered, because they are fortuitous records. The screening process included the removal of duplicates, title and abstract screening, and full-text assessment of the selected articles [23]. This approach allowed us to rigorously and transparently synthesize the available knowledge on the diet of post-metamorphic anurans in Brazil, identifying gaps and trends in the literature. We considered only studies published until June 30th, 2025 and included anuran species occurring within the territory of the Brazilian federation, based on the list of amphibians in Brazil [15], which compiled 1,144 valid species. Additionally, as this list was outdated at the time of our search, we incorporated taxonomic updates from Frost [14], including newly described species for Brazil and recent taxonomic rearrangements, which increased the number of Brazilian species to 1,232 spp. To ascertain the ecoregion distribution of each species, we referenced the ICMBio database [24] and studies on the distribution of species.

From each study, the following variables were extracted: (i) taxonomic classification (species, genus, family), (ii) method used to obtain the diet (dissection, stomach flush, field observation), (iii) assessment of prey availability, (iv) origin of sampled individuals (whether they were collected in the field or were from museum specimens) and, (v) source scientific journal and its 2022 Clarivate impact factor. In addition, for each species, we obtained (vi) the ecoregion of occurrence [24], (vii) IUCN Red List conservation status [17], and (viii) national conservation status (Ordinance No. 300 of the Ministry of the Environment, dated December 13, 2022). The species names and the year of description were recorded according to Frost [14]. The classification of species distribution by Brazilian ecoregion was carried out by consulting the specialized bibliography and Frost [14].

3 Results

We retrieved 221 scientific studies describing the diet of anurans species in Brazil, published between 1971 and June 30th 2025 (Fig. 1). The first diet study was published by the journal *Ciência e Cultura*, describing the diet of *Thoropa miliaris* (Spix, 1824) [25].




Fig. 1 Evolution in the descriptions of the diet of Brazilian anurans between 1971 and 2025

Notably, only nine additional studies on anuran diets were identified throughout the entire 20th century. However, with the dawn of the 21st century, we observed a growth in the number of studies (Fig. 1). We observed an increase in the anuran diet publication of articles between 2019 and 2023 (69 studies published; 31.2%).

3.1 Species with described diet

We documented dietary information for a total of 233 species of anurans, representing approximately 19% of Brazilian species. The family with the highest number of species with a described diet was Hylidae (84, 36.1%), followed by Leptodactylidae (55; 26.6%), Bufonidae (23; 9.9%), and Microhylidae (15; 6.4%). For the families Alsodidae and Centrolenidae, no studies on the diet of their species were found (Fig. 2).

The species with the highest number of diet studies were *Leptodactylus fuscus* ($N=14$) and *Physalaemus cuvieri* ($N=14$), followed by *Boana raniceps* ($N=12$), *L. macrosternum* ($N=11$), *Dendropsophus minutus* ($N=9$), *L. luctator* ($N=7$), *L. podicipinus* ($N=7$), *Rhinella diptycha* ($N=7$), *Adenomera hylaedactyla* ($N=6$), *P. cicada* ($N=6$), *A. andreae* ($N=5$), *Allobates femoralis* ($N=5$), *B. albopunctata* ($N=5$), *L. pentadactylus* ($N=5$), *L. troglodytes* ($N=5$), *R. granulosa* ($N=5$), *Scinax fuscomarginatus* ($N=5$), *S. x-signatus* ($N=5$), *P. albifrons* ($N=5$), and *Pleurodema diplolister* ($N=5$).

Among the 233 species with documented diets, 133 (57.1%) species are represented by a single dietary description, 43 (18.4%) have two dietary studies and 57 (24.5%) have more than two studies. Among the 221 retrieved studies, 180 used the dissection method to examine amphibian diets (81.4%), 40 (18.1%) used stomach flushing, and two (1%) used the observation method. Nine (4.1%) studies used specimens deposited in scientific collections, 176 (79.6%) killed individuals to assess the diet, 35 (15.8%) captured and released individuals after stomach flushing and only 34 (15.4%) studies provided data on the availability of prey in the environment.




Fig. 2 Total number of species with described diet from each of the Brazilian anuran families




Fig. 3 Number of Brazilian anuran species in each of the categories of threat of the IUCN red list. Where: LC=Least Concern, NE=Not Evaluated, DD=Data Deficient, NT=Near Threatened, VU=Vulnerable, EN=Endangered, CR=Critically Endangered

3.2 Threatened species

We recorded dietary information for 12 globally threatened species, encompassing two Critically Endangered (CR) species [*Adelophryne maranguapensis* and *Proceratophrys ararype*], five Endangered (EN) [*Brachycephalus garbeanus*, *Crossodactylus timbuhy*, *Cycloramphus brasiliensis*, *Hylodes nasus*, and *H. perere*], and five Vulnerable (VU) [*Brachycephalus brunneus*, *Chiasmocleis alagoana*, *Phyllodytes tuberculatus*, *Pseudopaludicola restinga*, and *Xenohyla truncata*] (Fig. 3). Regarding threatened species at national level, there is information on the diet only for *A. maranguapensis*, *C. alagoana*,




Fig. 4 Number of anuran species (overall and endemic) in each Brazilian ecoregion with one or more published diet descriptions

P. ararype, and *X. truncata* (Fig. 3). Except for *P. tuberculatus* (stomach-flushing method) and one population of *X. truncata* (observation method), all other threatened species were collected, killed, and dissected to obtain food contents.

None of the 62 species classified as Data Deficient (DD) at the international level have a described diet, and only one of the 29 species classified as Near Threatened (NT) has information about its diet (*Crossodactylus trachystomus*) (Fig. 3). Among the 103 Non-Evaluated (NE) species, only three species have available information on their diet (*Dendropsophus tapacurensis*, *Odontophrynus asper*, and *Pristimantis crepitaculus*), and regarding the 852 non-threatened species, classified as Least Concern (LC), 217 (25.5%) have a described diet (Fig. 3). At the national level, of the 85 species classified as DD, there is information on the diet only for *Phyllodytes tuberculatus*, while there is no information on the diet of any of the 27 species classified as NT (Fig. 3).

3.3 Diet analysis by ecoregion

The ecoregion with the highest number of dietary studies was the Atlantic Forest (110 publications; 49.8%), followed by the Amazon (47; 21.3%), Cerrado (32; 14.5%), Caatinga (23; 10.4%), Pampa (10; 4.5%) and Pantanal (8; 3.6%) (Fig. 4). We registered 91 endemic species considering all six Brazilian ecoregions with the described diet (10.1%) of a total of 907 endemic species (Table S1, Fig. 4). Of these species, 50 species are endemic to the Atlantic Forest (54.9%), 32 to the Amazon (35.2%), six to the Cerrado (6.6%), and three to the Caatinga (3.3%) (Table S1, Fig. 4).

3.4 Scientific journals

We identified 56 scientific journals that have published studies describing the diet of Brazilian post-metamorphic anurans. Of these, 19 (33.9%) do not have an Impact Factor (IF). Among the remaining journals, the average IF was 1.4 (± 1.1). *Iheringia (Zoology)* and *Phyllomedusa – Journal of Herpetology* had the lowest IF (0.4), while *Ecology* and *Journal of Animal Ecology* had the highest (4.8).

More specifically, 32.2% ($N=75$) of the 233 studies were published in journals without IF. Furthermore, the journal with the highest number of dietary studies published was *Herpetology Notes* ($N=26$; 11.2%), which also does not have IF. In contrast, the second journal with the highest number of publications was *North-Western Journal of Zoology* ($N=21$; 9%), which has an IF of 0.7. Regarding the other journals with the highest number of published studies, the *Brazilian Journal of Biology* ($N=16$; 6.9%) also does not have an IF value, while the *Journal of Herpetology* (13; 5.6%), *South American Journal of Herpetology* (13; 5.6%) and *Phyllomedusa – Journal of Herpetology* (10; 4.3%) have an IF value below one. The journals with the highest IF values that published studies on the diet of anurans were *Ecology* (IF = 4.8; $N=1$ study), *Journal of Animal Ecology* (IF = 4.8; $N=1$), *OIKOS* (IF = 3.4; $N=1$), *Oecologia* (IF = 2.7; $N=2$), *Copeia* (IF = 2.6; $N=3$), *Hydrobiologia* (IF = 2.6; $N=1$), *Herpetologica* (IF = 2.4; $N=1$) and *Journal of Zoology* (IF = 2; $N=1$), representing only 3.6% of published studies.

4 Discussion

Our review reveals striking gaps and biases in the current knowledge of the trophic ecology of Brazilian anurans. Diet data are still unavailable for the vast majority of species (> 80%), including species with wide geographic distributions and several that are threatened. This scarcity of basic ecological information as the feeding behavior, impose limits to our ability to understand species' functional roles in ecosystems, hinders the identification of key prey interactions, and constrains conservation planning. On the other hand, our findings underscore a broader pattern: while generalist species with large distributions are relatively well-studied, rare and range-restricted taxa—particularly those in under-sampled ecoregions such as the Caatinga, Pantanal, and Pampa—remain critically understudied. The recent increase in dietary studies is likely associated with the scientific funding crisis that affected Brazil in the late 2010s, rather than with increased academic incentives [26, 27]. Given their relatively low cost and logistical simplicity, dietary studies may have become a more accessible research alternative during periods of financial constraint. These knowledge gaps raise concerns not only for ecological research but also for evidence-based conservation strategies, especially as environmental changes accelerate across Brazil's diverse ecoregions.

4.1 Dietary research on Brazilian post-metamorphic anurans

Among the 1,232 known Brazilian anuran species, we identified 20 that have been more extensively studied, each with five or more published articles focusing on their trophic ecology: *Adenomera andreae* [28–32], *A. hylaedactyla* [28, 33–37], *Allobates femoralis* [28, 29, 32, 35, 38], *Boana albopunctata* [9, 32, 39–41], *B. raniceps* [9, 33, 34, 37, 42–48], *Dendropsophus minutus* [9, 29, 33, 34, 49–53], *Leptodactylus fuscus* [9, 30, 34, 37, 39, 43, 44, 54–60], *L. luctator* [9, 57, 61–65], *L. macrosternum* [9, 33, 34, 37, 43, 44, 59, 66–69], *L. pentadactylus* [30, 37, 70–72], *L. podicipinus* [9, 32, 37, 66, 73–75], *L. troglodytes* [33, 34, 43, 44, 76], *Physalaemus albifrons* [34, 43, 44, 77, 78], *P. cicada* [34, 43, 44, 77–79], *P. cuvieri* [9, 33, 34, 50, 56, 57, 74, 78, 80–85], *Rhinella diptycha* [9, 34, 43, 44, 56, 80, 86], *R. granulosa* [32, 34, 43, 44, 79], *Scinax fuscomarginatus* [9, 32, 34, 87, 88], and *S. x-signatus* [33, 34, 37, 43, 44]. These species are all classified as Least Concern (LC) and have broad geographic distributions. For example, *D. minutus* and *L. fuscus* are found

in all six Brazilian ecoregions [14, 17]. The wide distribution and high abundance likely explains the higher number of studies on these species.

Most of these species are generalists, with opportunistic feeding behavior [33, 34, 43–45, 56, 74]. However, this is not a rule, as *A. hylaedactyla*, *P. albifrons*, and *P. cuvieri* consume more ants than other insects—although studies on prey availability for these species were not evaluated—while *B. raniceps* has a preference for Coleoptera, and *P. cicada* is a termite specialist [79, 88]. Diet preference is a point in the natural history of species that must be known and understood in order to preserve the species. Another point of the diet that must be evaluated is ontogenetic differentiation. In fact, ontogenetic effects were noted, particularly in *D. minutus* and *L. podicipinus*, where juveniles consume smaller prey compared to adults [29, 73, 74]. The assessment of ontogenetic differences and food preferences is important when making conservation decisions, such as when evaluating endangered species.

Additionally, nearly 50% of studies on these species also analyzed community-level dietary patterns, examining niche differentiation, food resource partitioning, and intra-specific dietary variation [29, 33, 34, 43, 44, 81]. These studies typically demand substantial time and resources but provide insights into evolutionary, ecological, and historical influences on dietary patterns [e.g., 9, 10]. Such integrative approaches not only advance our understanding of trophic ecology but also tend to result in publications in higher-impact ecological journals, underscoring the scientific value of community-level dietary analyses.

Most anuran species with available dietary studies are considered generalists with opportunistic feeding behavior; however, a few exceptions have been documented. Three species exhibit specialist diets: *Amazophrynella minuta* appears to specialize in mites (Acari) [89], *Rhinella granulosa* in termites (Isoptera) [79], and *Phyllodytes edelmoi* in ants (Formicidae) [90]. In addition, several other species demonstrate clear dietary preferences. For instance, *Aplastodiscus albosignatus* and *Boana semiguttata* preferentially consume Coleoptera [91], *B. raniceps* favors Coleoptera and Hemiptera [47], *Lepidactylus notoaktites* feeds mainly on Hemiptera and Pseudoscorpionida [91], *Pseudis minuta* on Hemiptera, Coleoptera, Orthoptera, and Araneae [92], *Scinax acuminatus* on Hemiptera [47], *Physalaemus ephippifer* on Formicidae [60], *Proceratophrys boiei* on Orthoptera [91], and *Trachycephalus typhonius* on Araneae, Coleoptera and Hemiptera [47]. Moreover, *Physalaemus carrizorum* tends to consume primarily Formicidae [93], while *Scythrophrys sawayae* and *Brachycephalus brunneus* show a marked preference for Acari [91, 94]. Additionally, *Allobates marchesianus* and *Anomaloglossus stepheni* appear to have a preference for ants and Collembola [95]. These findings highlight that, although generalist feeding is predominant among Brazilian anurans, dietary specialization and preferences do occur and may reflect ecological adaptations, habitat use, or evolutionary history—underscoring the importance of expanding dietary studies to a broader range of species and ecological contexts.

4.2 Threatened species

Despite the large number of threatened species, both internationally (184 spp.; 14.9%) [17] and nationally (56 spp.; 4.5%; Ordinance No. 300, of December 13, 2022), only 15 (6.8%) studies describe the diet of 12 globally threatened species. Seven of the 12 threatened species with known diets were generalists, typical of amphibians, exhibiting

opportunistic feeding behaviors [78, 96–101]. In contrast, *B. brunneus* [94] and *B. garbeanus* [102] had a more specialized diet, primarily feeding on mites, suggesting active foraging behavior. Regarding foraging strategies, *A. maranguapensis* was identified as an active forager [80], while *C. timbuhy* exhibited both active and “sit-and-wait” foraging behaviors, feeding on both mobile insects (e.g. ants, flies) and stationary prey (e.g. insect larvae) [96]. *Crossodactylus brasiliensis* consumes both terrestrial and aquatic prey, feeding in and out of the water [99]. An exception to the general trend of invertebrate-based diets is *X. truncata*, which feeds actively on fruits, nectar, and plant parts [103–105]. Recently, *X. truncata* was observed feeding on nectar from *Iris x germanica* and *Cordia taguahyensis* flowers, with plant pollen adhering to its body, suggesting its role as a potential pollinator [105].

Regarding threat assessment, it is noteworthy that no dietary data are available for the 62 species (5.1%) currently classified as Data Deficient (DD) [17]. This is particularly concerning given that, with the exception of *Physalaemus claptoni*, all other DD species listed by the IUCN are endemic to Brazilian ecoregions—40 species (64%) are endemic to the Atlantic Forest, 16 (26%) to the Amazon, and five (8%) to the Cerrado [17; Ordinance No. 300, December 13, 2022]. Among them, the lack of dietary information for *Cycloramphus asper* is especially notable. In addition to being endemic to the Atlantic Forest, this species is listed as Near Threatened (NT) at the national level (Ordinance No. 300, December 13, 2022). These gaps underscore the urgent need for more research not only on the trophic ecology of these species, but also on their broader natural history, to better assess their actual conservation status.

This research highlights a significant knowledge gap regarding the feeding ecology of Brazil’s most threatened anuran species. The scarcity of studies in this area is likely due to the fact that many of these species have small, isolated populations and are often endemic [17], as is the case with DD species. We emphasize that dietary studies on threatened species are essential for understanding ecological interactions, identifying factors that influence survival and reproductive success, and evaluating species’ resilience to environmental changes [2–5]. Therefore, new approaches and increased sampling efforts are urgently needed to obtain reliable dietary data for the remaining threatened species. Furthermore, we propose that trophic information can serve as an early indicator of ecosystem changes or anthropogenic impacts, thereby informing habitat management strategies [106]. Integrating dietary analyses into conservation research is crucial for the development of effective management plans aimed at safeguarding threatened amphibian species.

4.3 Ecoregions and endemic species

Brazil harbors a high number of anuran species endemic to one of its six major ecoregions (i.e., Amazon, Atlantic Forest, Caatinga, Cerrado, Pampa, and Pantanal), totaling 907 species, and representing 73.6% of the country’s anuran fauna [24]. Despite this, only 91 of these species (10.1%) have dietary information available in the literature, highlighting a significant research gap and suggesting that endemic species remain disproportionately understudied compared to their non-endemic counterparts.

The Atlantic Forest (493 spp.; 54.3%) and the Amazon (318 spp.; 35.1%) are the ecoregions with the highest number of endemics. However, only 50 and 32 of them, respectively, have available dietary data. A similar pattern is observed in the Cerrado, which

hosts 68 endemic species (7.5%), yet only six have documented dietary information. In contrast, the Caatinga (19 spp.), Pampa (five spp.), and Pantanal (one spp.) are the ecoregions with the lowest numbers of endemic anurans, together accounting for approximately 3% of Brazil's endemic species. Among these, only three species endemic to the Caatinga have their diet described [96, 97].

This pronounced disparity between the number of endemic anuran species and the availability of dietary information reveals a critical gap in our understanding of Brazil's amphibian trophic ecology [107]. Endemic species, by definition, have restricted geographic distributions and are often more vulnerable to habitat loss, climate change, and other anthropogenic pressures [108, 109]. The lack of basic ecological data—such as dietary composition—not only hampers conservation planning but also limits our ability to understand functional roles and ecological interactions within their native ecosystems [107, 110, 111]. Addressing this knowledge gap is essential for developing effective conservation strategies and for advancing ecological theory in biodiverse and threatened ecoregions [110, 111].

4.4 Scientific journals

Among the 56 scientific journals publishing studies on the trophic ecology of Brazilian anurans, about 1/3 of the articles (33.9%) were published in journals without an impact factor. Meanwhile, most articles on the diet of Brazilian amphibians (44.8%) were published in scientific journals with an impact factor of less than one. This suggests that studies on trophic ecology tend to be published in journals with lower impact factors, which is concerning, as studies on the natural history of species may lack scientific prestige [112], with high-impact journals often avoiding publishing these studies [19]. This pattern is worrying because trophic ecology is vital for understanding habitat use and dietary preferences, which are key for experimental design and testing ecological hypotheses [1–5].

In contrast, only 11 studies (5%) were published in journals with an Impact Factor (IF) greater than two. Among them, two studies were published in *Ecology* and the *Journal of Animal Ecology* (IF = 4.8), addressing niche variation in dietary patterns [81] and spatiotemporal variation in predator-prey interactions [10]. Other studies in higher-impact journals typically combine trophic ecology with additional ecological factors and more robust analyses [29, 30, 33, 34, 113–115], or focus on endangered species, such as *Xenohyla truncata* [105]. Therefore, while studies solely describing species' diets are important, including other approaches and detailed statistical analyses increases the likelihood of publication in high-impact journals.

4.5 Methods in diet studies of Brazilian Anuran amphibians

Different methods can be used to assess the anurans diet. In the present study, we detected the use of stomach and intestine dissection, stomach flush, observation method and isotope analysis. The most of studies on the diet of Brazilian anuran amphibians (180 publications; 81.4%) used stomach and intestine dissection. Only nine studies (4.1%) analyzed museum specimens, thereby avoiding the need to collect and kill new animals.

In recent decades, there has been a notable increase in the use of stomach flushing—a non-lethal technique proposed by Solé et al. [116], which force individuals to regurgitate stomach contents without harm. Despite its ethical and conservation advantages,

the use of this method remains limited, appearing in only 40 publications (18.1%), with just one study applying it to a threatened species, *Phyllodytes tuberculatus* [117]. Given the growing number of threatened and data-deficient species, we strongly encourage the broader adoption of stomach flushing, particularly when working with populations that are small, declining, or difficult to access. Its implementation can significantly reduce the impact of fieldwork on vulnerable species and promote more sustainable research practices.

Additionally, two studies used the observation method to describe the trophic ecology of *Phantasmarana apuana* [118] and *Xenohyla truncata* [105], a method that, while logistically challenging, provides valuable data without harming individuals.

Some studies have also employed stable isotope analysis to understand anuran diets [64, 81, 93, 113, 114], offering insights into nutritional sources and dietary variation. Stable isotopes help identify whether a species' diet consists of terrestrial or aquatic prey [113, 114], whether isotopic niche breadth varies by season [96], and determine the origin of forest species [93]. While this technique is invaluable for understanding species' diets and their environmental adaptations, the costs of such analyses can be prohibitive for mid-income countries with limited science funding. Nonetheless, stable isotope analysis remains crucial for unraveling the complexities of trophic ecology and aiding in biodiversity conservation.

Furthermore, information on prey availability is crucial for detecting dietary selection [3, 6, 7, 11], and can show preferences and help us understand the requirements for species conservation. In this sense, only 34 studies (15.4%) included data on prey availability in the environment. Over half of these studies were published in the last decade [64, 75, 76, 91, 93, 119–122], indicating a growing trend. Assessing prey availability for arboreal species remains challenging, as most tree frogs forage in the canopy during non-reproductive periods, complicating prey assessments outside of their breeding sites (ponds, streams).

More recently, DNA-based techniques have been increasingly applied in trophic ecology studies [123]. Methods such as multiplex PCR and next-generation sequencing (NGS) have become powerful tools, enabling the identification of prey items with high sensitivity and taxonomic resolution, even when morphological identification is hindered by digestion. Despite their advantages—including the ability to analyze complex diets from minimal or highly degraded samples—these techniques are still rarely employed in Brazilian herpetology. This represents a promising opportunity to advance dietary research and improve our understanding of anuran trophic ecology in the region.

4.6 Information gaps and future directions

Despite the increasing number of studies on the trophic ecology of Brazilian anurans, significant gaps remain. Currently, we know the feeding ecology of fewer than 20% of species, with most species only studied in a single location, underscoring the need for further research in this area.

As new technologies and research tools, such as those in genetics and molecular biology, evolve, fewer researchers are pursuing studies on natural history [19, 112]. Research in natural history often involves labor-intensive, costly, and time-consuming fieldwork. Moreover, journals that publish natural history studies generally have a low scientific impact, which discourages researchers from investing in this type of research. This gap

is concerning as natural history knowledge contributes to all stages of scientific research, from hypothesis development to data interpretation and discussion [16, 19, 112]. While natural history studies do not typically require costly technologies, integrating such methods can improve publication chances [19]. For example, Huckembeck et al. [64, 113, 114], Araújo et al. [81], and De Oliveira et al. [93] used stable isotopes to infer the nutritional contributions of primary producers in studying anuran diets, leading to publications in high-impact journals compared to most diet-focused studies.

We particularly emphasize the need for more research in the Atlantic Forest, Amazon and Cerrado. Despite being the biomes with the most publications, the biodiversity in these regions still requires substantial efforts to evaluate species' diets. This information gap extends to the Caatinga, Pampa, and Pantanal, the least-studied biomes regarding amphibian trophic ecology. Notably, only three endemic species from the Caatinga, *Adelophryne maranguapensis*, *Adelophryne baturitensis*, and *Proceratophrys ararype*, have their diets studied.

While some species have been identified as conservation priorities, the current lack of knowledge about the trophic ecology of Brazilian anurans signals the need for more studies across practically all species. This review emphasizes the critical need for more research on the diets of anurans, as such data are essential for understanding predator-prey interactions, hypothesis testing, and assessing conservation status. We recommend fieldwork using non-lethal methods that include as many species as possible. Studies on specimens in scientific collections are also vital to address this information gap.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1007/s44396-025-00008-7>.

Supplementary Material 1.

Acknowledgements

KC, DJS, MS and AMT thank CNPq for research fellowships (CNPq, process number 444240/2024-1, 311284/2023-0, 309365/2019-8, and 305533/2022-0, respectively). MS also acknowledges a CAPES/Alexander von Humboldt Foundation grant (BEX 0585/16–5).

Author contributions

C.F.M. collected the data. C.F.M. e K.C. wrote the main manuscript text. K.C. prepared figures. C.F.M., K.C., L.K.S., D.J.S., S.I., and A.M.T. reviewed the manuscript.

Funding

No funding was received to assist with the preparation of this manuscript.

Data availability

No datasets were generated or analysed during the current study.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Instituto de Biologia, Universidade Estadual de Campinas (UNICAMP), Campinas, São Paulo, Brazil

²Laboratório de Interações Ecológicas e Biodiversidade (LIEB), Departamento de Biologia, Universidade Federal do Ceará (UFC), Fortaleza, Ceará, Brazil

³Department of Biology, The Pennsylvania State University, University Park, United States of America

⁴Instituto de Biociências, Universidade Federal de Mato Grosso do Sul (UFMS), Campo Grande, Mato Grosso do Sul, Brazil

⁵Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz (UESC), Ilhéus, Bahia, Brazil

⁶Museum Koenig Bonn (ZFMK), Leibniz Institute for the Analysis of Biodiversity Change, Bonn, Germany

⁷Universidade do Oeste de Santa Catarina (UNOESC), Joaçaba, Santa Catarina, Brazil

⁸Laboratório de Ecologia de Vertebrados Terrestres (LEVERT), Universidade do Vale do Rio dos Sinos (UNISINOS), Porto Alegre, Rio Grande do Sul, Brazil

⁹Instituto Biodiversa, Porto Alegre, Rio Grande do Sul, Brazil

Received: 10 March 2025 Accepted: 10 July 2025

Published online: 13 August 2025

References

1. Anderson J. Why ecology needs natural history. *Am Sci.* 2017;105: 290. <https://doi.org/10.1511/2017.105.5.290>.
2. Pianka ER. The structure of lizard communities. *Annu Rev Ecol Syst.* 1973;4:53–74. <https://doi.org/10.1146/annurev.es.04.11.0173.000413>.
3. Duellman WE, Trueb L. *Biology of amphibians.* JHU; 1994.
4. Sih A, Christensen B. Optimal diet theory: when does it work, and when and why does it fail? *Anim Behav.* 2001;61:379–90. <https://doi.org/10.1006/anbe.2000.1592>.
5. Solé M, Rödder D. Dietary assessments of adult amphibians. *Amphibian ecology and conservation: a handbook of techniques.* Oxford: Oxford University Press; 2010. pp. 167–84.
6. Toft CA. Feeding ecology of thirteen syntopic species of anurans in a seasonal tropical environment. *Oecologia.* 1980;45:131–41. <https://doi.org/10.1007/BF00346717>.
7. Toft CA. Feeding ecology of Panamanian litter anurans: patterns in diet and foraging mode. *J Herpetol.* 1981;15:139. <https://doi.org/10.2307/1563372>.
8. Estes JA, Terborgh J, Brashares JS, Power ME, Berger J, Bond WJ, Carpenter SR, Essington TE, Holt RD, Jackson JBC, Marquis RJ, Oksanen L, Oksanen T, Paine RT, Pikitch EK, Ripple WJ, Sandin SA, Scheffer M, Schoener TW, Shurin JB, Sinclair ARE, Soulé ME, Virtanen R, Wardle DA. Trophic downgrading of planet Earth. *Science.* 2011;333:301–6. <https://doi.org/10.1126/science.1205106>.
9. Ceron K, Provete DB, Pires MM, Araujo AC, Blüthgen N, Santana DJ. Influence of prey availability in the dissimilarity of predator–prey interactions. *Bull Ecol Soc Am.* 2022;103: e01989. <https://doi.org/10.1002/bes2.1989>.
10. Ceron K, Provete DB, Pires MM, Araujo AC, Blüthgen N, Santana DJ. Differences in prey availability across space and time lead to interaction rewiring and reshape a predator–prey metaweb. *Ecology.* 2022;103:e3716. <https://doi.org/10.1002/ecy.3716>.
11. Anderson AM, Hauko DA, Anderson JT. Diet composition of three anurans from the Playa wetlands of Northwest Texas. *Copeia.* 1999;1999:515. <https://doi.org/10.2307/1447502>.
12. Schoener TW. Resource partitioning in ecological communities: research on how similar species divide resources helps reveal the natural regulation of species diversity. *Science.* 1974;185:27–39. <https://doi.org/10.1126/science.185.4145.27>.
13. Hortal J, De Bello F, Diniz-Filho JAF, Lewinsohn TM, Lobo JM, Ladle RJ. Seven shortfalls that beset large-scale knowledge of biodiversity. *Annu Rev Ecol Syst.* 2015;46:523–49. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>.
14. Frost DR. (2024) *Amphibian Species of the World: an Online Reference.* Version 6.2.
15. Segalla MV, Berneck B, Canedo CC, Grant T, Caramaschi U, Cruz CAG, Garcia PCA, Haddad CFB, Lourença AC, Mângia S, Mott T, Nascimento LB, Toledo LF, Werneck FP, Langone JA. List of Brazilian amphibians. *Herpetologia Brasileira.* 2021;10:121–216. <https://doi.org/10.5281/ZENODO.4716176>.
16. Guerra V, Llusia D, Gambale PG, Morais ARD, Márquez R, Bastos RP. The advertisement calls of Brazilian anurans: historical review, current knowledge and future directions. *PLoS One.* 2018;13: e0191691. <https://doi.org/10.1371/journal.pone.0191691>.
17. IUCN. (2024) *The IUCN Red List of Threatened Species.* Version 2024-1.
18. Luedtke JA, Chanson J, Neam K, Hobin L, Maciel AO, Catenazzi A, Borzée A, Hamidy A, Aowphol A, Jean A, Sosa-Bartuano Á, Fong GA, De Silva A, Fouquet A, Angulo A, Kidov AA, Muñoz Saravia A, Diesmos AC, Tominaga A, Shrestha B, Gratwicke B, Tjaturadi B, Martínez Rivera CC, Vásquez Almazán CR, Señaris C, Chandramouli SR, Strüssmann C, Cortez Fernández CF, Azat C, Hoskin CJ, Hilton-Taylor C, Whyte DL, Gower DJ, Olson DH, Cisneros-Heredia DF, Santana DJ, Nagombi E, Najafi-Majid E, Quah ESH, Bolaños F, Xie F, Brusquetti F, Álvarez FS, Andreone F, Glaw F, Castañeda FE, Kraus F, Parra-Olea G, Chaves G, Medina-Rangel GF, González-Durán G, Ortega-Andrade HM, Machado IF, Das I, Dias IR, Urbina-Cardona JN, Crnobrnja-Isailović J, Yang J-H, Jianping J, Wangyal JT, Rowley JJJ, Measey J, Vasudevan K, Chan KO, Gururaja KV, Ovaska K, Warr LC, Canseco-Márquez L, Toledo LF, Díaz LM, Khan MMH, Meegaskumbura M, Acevedo ME, Napoli MF, Ponce MA, Vaira M, Lampo M, Yáñez-Muñoz MH, Scherz MD, Rödel M-O, Matsui M, Fildor M, Kusriini MD, Ahmed MF, Rais M, Kouamé NG, García N, Gonwouo NL, Burrows PA, Imbun PY, Wagner P, Kok PJR, Joglar RL, Auguste RJ, Brandão RA, Ibáñez R, Von May R, Hedges SB, Biju SD, Ganesh SR, Wren S, Das S, Flechas SV, Ashpole SL, Robleto-Hernández SJ, Loader SP, Incháustegui SJ, Garg S, Phimmachak S, Richards SJ, Slimani T, Osborne-Naikatini T, Abreu-Jardim PTF, Condez TH, De Carvalho TR, Cutajar TP, Pierson TW, Nguyen TQ, Kaya U, Yuan Z, Long B, Langhammer P, Stuart SN. (2023) Ongoing declines for the world's amphibians in the face of emerging threats. *Nature.* 622:308–314. <https://doi.org/10.1038/s41586-023-06578-4>
19. Mccallum ML, Mccallum JL. Publication trends of natural history and field studies in herpetology. *Herpetol Conserv Biol.* 2006;1:62–7.
20. Nunes-de-Almeida C. A revised classification of the amphibian reproductive modes. *Salamandra.* 2021;57:413–27.
21. Silla AJ, Calatayud NE, Trudeau VL. Amphibian reproductive technologies: approaches and welfare considerations. *Conserv Physiol.* 2021;9: coab011. <https://doi.org/10.1093/conphys/coab011>.
22. Méndez-Tepepa M, Morales-Cruz C, García-Nieto E, Anaya-Hernández A. A review of the reproductive system in Anuran amphibians. *Zoological Lett.* 2023;9:3. <https://doi.org/10.1186/s40851-023-00201-0>.
23. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Aki EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ n71.* 2021. <https://doi.org/10.1136/bmj.n71>.

24. ICMBio. Sistema de Avaliação do Risco de Extinção da Biodiversidade – SALVE; 2024.
25. Sazima I. The occurrence of marine invertebrates in the stomach contents of the frog *Thoropa miliaris*. *Cienc Cult*. 1971;23:647–8.
26. Hipólito J, Shirai LT, Diele-Viegas LM, Halinski R, Soares Pires CS, Fontes EMG. Brazilian budget cuts further threaten gender equality in research. *Nat Ecol Evol*. 2021;6:234–234. <https://doi.org/10.1038/s41559-021-01640-8>.
27. Knockaert M, Deschryvere M, Lecluyse L. The relationship between organizational interdependence and additionality obtained from innovation ecosystem participation. *Sci Public Policy*. 2019;46:490–503. <https://doi.org/10.1093/scipol/scz002>.
28. Vitt L, Caldwell JP. Resource utilization and guild structure of small vertebrates in the Amazon forest leaf litter. *J Zool*. 1994;234:463–76. <https://doi.org/10.1111/j.1469-7998.1994.tb04860.x>.
29. Lima A, Magnusson WE. Partitioning seasonal time: interactions among size, foraging activity and diet in leaf-litter frogs. *Oecologia*. 1998;116:259–66. <https://doi.org/10.1007/s004420050587>.
30. Caldwell JP, Vitt LJ. Dietary asymmetry in leaf litter frogs and lizards in a transitional Northern Amazonian rain forest. *Oikos*. 1999;84:383. <https://doi.org/10.2307/3546419>.
31. Sabagh LT, Mello RDS, Rocha CFD. Food niche overlap between two sympatric leaf-litter frog species from central Amazonia. *Zoologia (Curitiba)*. 2012;29:95–8. <https://doi.org/10.1590/S1984-46702012000100013>.
32. Ferreira-Silva SK, Alexandre RJR, Pena SA, Lucena MDL, Vieira TB, Gomes FBR. Associations between morphological attributes and food resources in anurans from the middle Xingu region, Brazil. *Braz J Biol*. 2025;85:1–13. <https://doi.org/10.1590/01519-6984.274592>.
33. Leite-Filho E, De Oliveira FA, Eloi FJ, Liberal CN, Lopes AO, Mesquita DO. Evolutionary and ecological factors influencing an anuran community structure in an Atlantic rainforest urban fragment. *Copeia*. 2017;105:64–74. <https://doi.org/10.1643/C-H-15-298>.
34. Caldas FLS, Garda AA, Cavalcanti LBQ, Leite-Filho E, Faria RG, Mesquita DO. Spatial and trophic structure of anuran assemblages in environments with different seasonal regimes in the Brazilian Northeast region. *Copeia*. 2019;107:567. <https://doi.org/10.1643/CH-18-109>.
35. Almeida BC, Santos RS, Santos TF dos, de Souza MB, Menin M. (2019) Diet of five anuran species in a forest remnant in eastern Acre state, Brazilian Amazonia. *Herpetol Notes* 12:945–952.
36. Sanches PR, Pedrosa-Santos F, Costa-Campos CE. Diet of *Adenomera hylaedactyla* (Cope, 1868) (Anura: Leptodactylidae) from an urban area in Southern Amapá, Eastern Amazon. *Herpetol Notes*. 2019;12:841–5.
37. Costa-Campos CE, Sales RFD, Freire EMX. Trophic ecology of an assemblage of anurans from the Amazonian savanna in Amapá, Brazil. *North-West J Zool*. 2024;20:151–5.
38. Simões PI, Lima AP, Farias IP. The description of a cryptic species related to the pan-Amazonian frog *Allobates femoralis* (Boulenger 1883) (Anura: Aromobatidae). *Zootaxa*. 2010;2406:1. <https://doi.org/10.11646/zootaxa.2406.1.1>.
39. Araújo MS, Dos Reis SF, Giaretta AA, Machado G, Bolnick DI. (2007) Intrapopulation diet variation in four frogs (Leptodactylidae) of the Brazilian Savannah. *Copeia* 2007:855–865. [https://doi.org/10.1643/0045-8511\(2007\)7\[855:IDVIF\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2007)7[855:IDVIF]2.0.CO;2).
40. Guimarães TCS, De Figueiredo GB, Mesquita DO, Vasconcelos MM. Ecology of *Hypsiboas albopunctatus* (Anura: Hylidae) in a Neotropical savanna. *J Herpetol*. 2011;45:244–50. <https://doi.org/10.1670/09-255.1>.
41. Pacheco EO, Ferreira VG, Carvalho RMHD. Diet of *Boana albopunctata* (Anura: Hylidae) in an Atlantic forest fragment of southeastern Brazil. *Phyllomedusa*. 2017;16:57–62. <https://doi.org/10.11606/issn.2316-9079.v16i1p57-62>.
42. Lt S, Vi F, Rocha Cfd. Living together, sometimes feeding in a similar way: the case of the syntopic hylid frogs *Hypsiboas raniceps* and *Scinax acuminatus* (Anura: Hylidae) in the Pantanal of Miranda, Mato Grosso do Sul state, Brazil. *Braz J Biol*. 2010;70:955–9. <https://doi.org/10.1590/S1519-69842010000500006>.
43. Protázio AS, Albuquerque RL, Falkenberg LM, Mesquita DO. Niche differentiation of an anuran assemblage in temporary ponds in the Brazilian semi-arid caatinga: influence of ecological and historical factors. *Herpetol J*. 2015;25:109–21.
44. Leite-Filho E, Vieira WLDS, Santana GG, Eloi FJ, Mesquita DO. Structure of a Caatinga anuran assemblage in Northeastern Brazil. *Neotrop Biol Conserv*. 2015;10:63–73. <https://doi.org/10.4013/nbc.2015.102.02>.
45. Sant'Anna BS, Fonseca AM, Ferreira RFB. Diet composition and coexistence of *Boana geographica* and *Boana raniceps* (Anura: Hylidae) from central Amazonia, Brazil. *North-West J Zool*. 2022;18:170–83.
46. Rodrigues NL, Castro Araújo K, Ávila RW, De Andrade EB. Feeding habits of *Boana raniceps* (Cope, 1862) in three mountains of Brazilian semi-arid. *Univ Sci*. 2023;28:257–77. <https://doi.org/10.11144/Javeriana.SC282.fhob>.
47. Machado R, Fagundes Cabral AA, Lima Serafim L, Cavalheri DG, Rodrigues Nehemy IK, Ceron K, Santana DJ. Trophic ecology of a Pantanal treefrog assemblage during a severe drought. *Stud Neotropical Fauna Environ*. 2025;1–10. <https://doi.org/10.1080/01650521.2025.2507368>.
48. Machado HTDS, Silva CFD, Benício RA, Ávila RW. Feeding ecology, reproductive biology, and sexual dimorphism of *Boana raniceps* (Anura: Hylidae) in an area of caatinga, Northeastern Brazil. *Caldasia*. 2023;46:71–80. <https://doi.org/10.15446/caldasia.v46n1.99220>.
49. Van Sluys M, Rocha CFD. Feeding habits and microhabitat utilization by two syntopic Brazilian Amazonian frogs (*Hyla Minuta* and *Pseudopaludicola* sp. (gr. *falcipes*)). *Braz J Biol*. 1998;58:559–62. <https://doi.org/10.1590/S0034-7108199800040003>.
50. Santos EM, Almeida AV, Vasconcelos SD. Feeding habits of six anuran (Amphibia: Anura) species in a rainforest fragment in northeastern Brazil. *Iheringia Ser Zool*. 2004;94:433–8. <https://doi.org/10.1590/S0073-47212004000400014>.
51. Solé M, Pelz B. Do male tree frogs feed during the breeding season? Stomach flushing of five syntopic hylid species in Rio Grande do Sul, Brazil. *J Nat Hist*. 2007;41:2757–63. <https://doi.org/10.1080/00222930701661282>.
52. Leivas PT, Mayer TB, Leivas FWT, Fávoro LF. Trophic niche of *Dendropsophus Minutus* (Anura: Hylidae) in Southern Brazil. *Phyllomedusa*. 2018;17:267–72. <https://doi.org/10.11606/issn.2316-9079.v17i2p267-272>.
53. Soeiro ALS, Guerreiro AKM, Oliveira-Souza AE, Santana MS, Costa-Campos CE. Diet composition of two hylid species (*Dendropsophus Haraldischultzi* and *D. minutus*) in anthropogenic environments in Eastern Amazonia. *Cuad Herpetol*. 2022;36:105–9. [https://doi.org/10.31017/CdH.2022.\(2021-067\)](https://doi.org/10.31017/CdH.2022.(2021-067)).
54. De-Carvalho CB, Freitas EBD, Faria RG, Batista RDC, Batista CDC, Coelho WA, Bocchiglieri A. História natural de *Leptodactylus Mystacinus* e *Leptodactylus fuscus* (Anura: Leptodactylidae) no Cerrado do Brasil central. *Biota Neotrop*. 2008;8:105–15. <http://doi.org/10.1590/S1676-06032008000300010>.

55. Sugai JLMM, Terra JDS, Ferreira VL. Diet of *Leptodactylus fuscus* (Amphibia: anura: Leptodactylidae) in the Pantanal of Miranda river, Brazil. *Biotà Neotrop.* 2012;12:99–104. <https://doi.org/10.1590/S1676-06032012000100008>.
56. Menin M, Santos RS, Borges RE, Piatti L. Notes on the diet of seven terrestrial frogs in three agroecosystems and forest remnants in Northwestern São Paulo state, Brazil. *Herpetol Notes.* 2015;8:401–5.
57. Marques-Pinto T, Barreto-Lima AF, Brandão RA. Dietary resource use by an assemblage of terrestrial frogs from the Brazilian Cerrado. *North-West J Zool.* 2019;15:135–46.
58. Santana DJ, Ferreira VG, Crestani GN, Neves MO. Diet of the rufous frog *Leptodactylus fuscus* (Anura, Leptodactylidae) from two contrasting environments. *Herpetozoa.* 2019;32:1–6. <https://doi.org/10.3897/herpetozoa.32.e35623>.
59. Baia RR, de Sanches J, Santos PR, Florentino FP, Costa-Campos AC. CE (2020) Diet overlap of three sympatric species of *Leptodactylus* Fitzinger (Anura Leptodactylidae) in a Protected area in the Brazilian Amazon. *Cuad Herpetol* 34:175–184. [https://doi.org/10.31017/CdH.2020.\(2020-006\)](https://doi.org/10.31017/CdH.2020.(2020-006)).
60. Silva IBD, Teixeira CC, Pinho SRC, Correa FS, Santos-Costa MCD. Prey availability in a forest fragment in Eastern Amazon: how is it related to the diet of anurans *Leptodactylus fuscus* and *Physalaemus ephippifer*? *An Acad Bras Ciênc.* 2025. <https://doi.org/10.1590/0001-3765202520241293>.
61. França L, Facure K, Giaretta A. Trophic and spatial niches of two large-sized species of *Leptodactylus* (Anura) in southeastern Brazil. *Stud Neotrop Fauna Environ.* 2004;39:243–8. <https://doi.org/10.1080/01650520400007330>.
62. Pazinato DMM, de O Trindade A, Oliveira SV, Cappellari LH. Dieta de leptodactylus latrans (Steffen, 1815) Na Serra do sudeste, Rio Grande do sul, Brasil. *Biotemas.* 2011;24:147–51.
63. De Oliveira M, Dalzochio MS, Da Silva Santos NLP, Tozetti AM. Prey selection by anurans in subtemperate swamps of the extreme south of Brazil. *South Am J Herpetol.* 2019;14:204. <https://doi.org/10.2994/SAJH-D-17-0005>.
64. Huckembeck S, Winemiller KO, Loebmann D, Garcia AM. Trophic structure of frog assemblages in coastal habitats in southern Brazil. *Austral Ecol.* 2020;45:977–89. <https://doi.org/10.1111/aec.12920>.
65. Mendonça Da Silva CM, Dalmolin† DA, Schuck LK, Moser CF, Tozetti AM. Predator morphology affects prey consumption: evidence from an Anuran population in subtropical wetlands. *HJ.* 2024;34:137–44. <https://doi.org/10.33256/34.3.137144>.
66. Piatti L, Souza F. Diet and resource partitioning among anurans in irrigated rice fields in Pantanal, Brazil. *Braz J Biol.* 2011;71:653–61. <https://doi.org/10.1590/S1519-69842011000400009>.
67. Costa DFSD, Oliveira JCDD, Oliveira JFD, Chaves MF, Silva JND, Sousa TPD. Dieta de *Leptodactylus macrosternum* (Amphibia; anura; Leptodactylidae) no Sertão Da paraíba, Brasil. *Revista Verde.* 2016;11:123. <https://doi.org/10.18378/rvads.v11i4.4372>.
68. Camurugi F, Magalhães F, de Henrique M, de Queiroz M C. Reproduction, sexual dimorphism, and diet of *Leptodactylus chaquensis* (Anura, Leptodactylidae) in Northeastern Brazil. *Herpetol Conserv Biol.* 2017;12:498–508.
69. Teles DA, Rodrigues JK, Teixeira AAM, de Sousa JA, Ribeiro JGG SC. Diet of *Leptodactylus macrosternum* (Miranda-Ribeiro 1926) (Anura: Leptodactylidae) in the Caatinga domain, Northeastern Brazil, Neotropical region. *Herpetol Notes.* 2018;11:223–6.
70. Galatti U. Population biology of the frog *Leptodactylus Pentadactylus* in a central Amazonian rainforest. *J Herpetol.* 1992;26:23. <https://doi.org/10.2307/1565017>.
71. Couto AP, Da Silveira R, Soares AVMV, Menin M. Diet of the smoky jungle frog *Leptodactylus Pentadactylus* (Anura, Leptodactylidae) in an urban forest fragment and in a pristine forest in central Amazonia, Brazil. *Herpetol Notes.* 2018;11:519–25.
72. Santos-Souza CJ, Rebelo-Silva K, Barbosa-Ferreira ME, Oliveira-Souza AE, Costa Anaissi JS, Sanches PR, Costa-Campos CE. Diet assessment of *Leptodactylus pentadactylus* (Anura, Leptodactylidae) from forest and savanna habitats in the Amazonian domain. *North-West J Zool.* 2022;18:184–9.
73. Rodrigues D, Uetanabaro M, Prado CPA. Seasonal and ontogenetic variation in diet composition of *Leptodactylus podicipinus* (Anura, Leptodactylidae) in the Southern Pantanal, Brazil. *Rev Esp Herpetol.* 2004;18:19–28.
74. Da Silva FR, Rossa-Feres DDC. Diet of anurans captured in forest remnants in southeastern Brazil. *Rev Esp Herpetol.* 2010;24:5–17.
75. Ceron K, De Toledo Moroti M, Alves Benício R, Porcel Balboa Z, Marçola Y, Becker Pereira L, Santana DJ. Diet and first report of batracophagy in *Leptodactylus podicipinus* (Anura: Leptodactylidae). *Neotrop Biodivers.* 2018;4:70–4. <https://doi.org/10.1080/23766808.2018.1467173>.
76. Oliveira MDS, Rodrigues NLA, Araújo KC, Andrade EB. Eating on the city: diet composition of *leptodactylus troglodytes lutz*, 1926 in Pedro II municipality, Piauí, Brazil. *Ensaios Ciênc.* 2023v;27:161–6. <https://doi.org/10.17921/1415-6938.2023v27n2p161-166>.
77. Protázio Ados, Protázio S, dos A, Mesquita S DO. Niche partitioning between two *Physalaemus* species (Anura, Leptodactylidae) in semiarid Caatinga in Northeast Brazil. *North-West J Zool.* 2019;15:157–67.
78. Oliveira CR, Cavalcante LA, Machado HTDS, Ávila RW, Morais DH. Diet of three sympatric *Physalaemus* (Anura: Leptodactylidae) from the Brazilian semiarid region. *Bol Mus Biol Mello Leitão.* 2021;16:13–9. <https://doi.org/10.46357/bcnaturais.v16i1.311>.
79. Santana As Juncá Fa. Diet of *Physalaemus cf. cicada* (Leptodactylidae) and *Bufo granulosus* (Bufonidae) in a semideciduous forest. *Braz J Biol.* 2007;67:125–31. <https://doi.org/10.1590/S1519-69842007000100017>.
80. Moreira G, Barreto L. Alimentação e Variação sazonal Na frequência de Capturas de Anuros Em Duas localidades do Brasil central. *Braz J Biol.* 1996;13:313–20. <https://doi.org/10.1590/S0101-81751996000200002>.
81. Araújo MS, Bolnick DI, Martinelli LA, Giaretta AA, Dos Reis SF. Individual-level diet variation in four species of Brazilian frogs. *J Anim Ecol.* 2009;78(4):848–56. <https://doi.org/10.1111/j.1365-2656.2009.01546.x>.
82. Leivas PT, Leivas FWT, Campião K. Diet and parasites of the Anuran *Physalaemus cuvieri* fitzinger, 1826 (Leiuperidae) from an Atlantic forest fragment. *Herpetol Notes.* 2018;11:109–13.
83. Gambale PG, Da Silva MR, Oda FH, Bastos RP. Diet and trophic niche of two sympatric *Physalaemus* species in central Brazil. *South Am J Herpetol.* 2020;17:63. <https://doi.org/10.2994/SAJH-D-17-00100.1>.
84. Braga HSN, Vieira MVSA, Silva TAF, Protázio AS, Protázio AS. Acoustic partitioning explains the coexistence between two *Physalaemus* species (Anura, Leptodactylidae) in the Atlantic forest in Eastern Bahia state, Brazil. *An Acad Bras Ciênc.* 2023;95: e20211348. <https://doi.org/10.1590/0001-3765202320211348>.
85. Farina RK, Moser CF, Scali S, Oliveira MD, Witt P, Tozetti AM. Diet and trophic niche overlap of four syntopic species of *Physalaemus* (Anura: Leptodactylidae) in Southern Brazil. *Acta Herpetol.* 2023;18:1–19. https://doi.org/10.36253/a_h-11294.
86. Batista R, de C, De-Carvalho CB, de Freitas EB, Batista C, de Coelho C, Faria WA RG. Diet of *Rhinella Schneideri* (Werner, 1894) (Anura: Bufonidae) in the cerrado, central Brazil. *Herpetol Notes.* 2011;4:017–21.

87. Michelin G, Ceron K, Santana DJ. Prey availability influences the diet of *Scinax fuscomarginatus* in a Cerrado area, central Brazil. *Anim Biodiv Conserv*. 2020;169–75. <https://doi.org/10.32800/abc.2020.43.0169>.
88. Ceron K, Paula T, Scheibler P, Fadel R, Guimarães CDS, Silva LA, Santana DJ. Trophic ecology of small to large hylids from an Amazonia-Cerrado transitional zone in Brazil. *Food Webs*. 2023;36: e00295. <https://doi.org/10.1016/j.fooweb.2023.e00295>.
89. Lima AP. The effects of size on the diets of six sympatric species of postmetamorphic litter anurans in central Amazonia. *J Herpetol*. 1998;32:392. <https://doi.org/10.2307/1565453>.
90. Araujo-Vieyra K, Gonçalves U, Guedes Dos Santos J, Gomes Ferreira T, Skuk G O. Feeding habits of the bromeligenous treefrog phyllodytes edelmoi peixoto, Caramaschi & freire, 2003 (Anura: Hylidae) from the state of Alagoas, Northeastern Brazil. *Cuad Herpetol*. 2018;32. [https://doi.org/10.31017/CdH.2018.\(2017-19\)](https://doi.org/10.31017/CdH.2018.(2017-19)).
91. Moser CF, Farina RK, Dudczak AC, Tozetti AM, Lingnau R. Feeding ecology of endemic frogs of the Atlantic forest in southern Brazil. *An Acad Bras Cienc*. 2022;94: e20210282. <https://doi.org/10.1590/0001-376520220210282>.
92. Farina RK, Moser CF, Schuck LK, Tozetti AM. Do human-induced habitat changes affect the feeding behaviour of frogs? The case of *Pseudis minuta* Günther, 1858. *Herpetol Notes*. 2024;17:161–7.
93. De Oliveira M, Moser CF, Rebelato MM, Camargo PB, Tozetti AM. Trophic ecology of two amphibian species in patches and core forest of Atlantic forest: A dietary and isotopic approach. *Austral Ecol*. 2021;47:278–90. <https://doi.org/10.1111/aec.13107>.
94. Fontoura PL, Ribeiro LF, Pie MR. Diet of *Brachycephalus brunneus* (Anura: brachycephalidae): in the Atlantic Rainforest of Paraná, southern Brazil. *Zoologia (Curitiba)*. 2011;28:687–9. <https://doi.org/10.1590/S1984-46702011000500019>.
95. Juncá F, Eterovick P. (2007) Feeding Ecology of two Sympatric Species of Aromobatidae, *Allobates marchesianus* and *Anomaloglossus stepheni*, in Central Amazon. *J Herpetol* 41:301–308. [https://doi.org/10.1670/0022-1511\(2007\)41\[301:FEOTSS\]2.0.CO;2](https://doi.org/10.1670/0022-1511(2007)41[301:FEOTSS]2.0.CO;2).
96. Araújo KC, Cassiano-Lima D, Brasileiro AC, Bezerra CH, Ávila RW. What the minute shield frogs *Adelophryne* eat at a humid forest relict in Ceará state, Northeastern Brazil. *North-West J Zool*. 2023;19:1–7.
97. Mascarenhas W, Oliveira CR, Ávila RW, Ribeiro SC. Diet and habitat use of *Proceratophrys ararype* (Anura: Odontophrynidae), an endemic frog from the Chapada do Araripe, Northeastern Brazil. *South Am J Herpetol*. 2023;26:56–62. <https://doi.org/10.2994/SAJH-D-21-00035.1>.
98. Campinhos EC, Ferreira RB, Mageski MM, Srbek-Araujo AC. Diet of *Crossodactylus timbuhy* (Anura: Hylodidae) in the reserva biológica Augusto Ruschi, state of Espírito Santo, Brazil. *Biota Neotrop*. 2020;20:e20190943. <https://doi.org/10.1590/1676-0611-bn-2019-0943>.
99. Maia-Carneiro T, Dorigo TA, Almeida-Gomes M, Van Sluys M, Rocha CFD. Feeding habits, microhabitat use, and daily activity of *Cycloramphus brasiliensis* (Anura: Cycloramphidae) from the Atlantic rainforest, Brazil. *Zoologia (Curitiba)*. 2012;29:277–9. <https://doi.org/10.1590/S1984-46702012000300007>.
100. Almeida VG, Yves A, Lima LMC, Novato TS, Neves MO. Natural history of *hylodes perere* Silva & Benmamoun, 2008 (Anura, Hylodidae) in the Serra Negra da Mantiqueira, Atlantic forest of Brazil: microhabitat, sexual dimorphism, diet and distribution. *J Nat Hist*. 2021;55:1477–90. <https://doi.org/10.1080/00222933.2021.1949065>.
101. Machado ADO, Winck G, Dorigo TA, Rocha CFD. Diet, diel activity pattern, habitat use, and reproductive effort of *Hylodes nasus* (Anura: Hylodidae) in one of the world's largest urban parks (Tijuca National Park), southeastern Brazil. *South Am J Herpetol*. 2016;11:127–35. <https://doi.org/10.2994/SAJH-D-16-00004.1>.
102. Dorigo TA, Siqueira CC, Vrcibradic D, Maia-Carneiro T, Almeida-Santos M, Rocha CFD. Ecological aspects of the pumpkin toadlet, *Brachycephalus garbeanus* Miranda-Ribeiro, 1920 (Anura: neobatrachia: Brachycephalidae), in a highland forest of southeastern Brazil. *J Nat Hist*. 2012;46:2497–507. <https://doi.org/10.1080/00222933.2012.713525>.
103. Da Silva HR, De Britto-Pereira MC. How much fruit do fruit-eating frogs eat? An investigation on the diet of *Xenohyla truncata* (Lissamphibia: anura: Hylidae). *J Zool*. 2006;270:692–8. <https://doi.org/10.1111/j.1469-7998.2006.00192.x>.
104. Da Silva HR, De Britto-Pereira MC, Caramaschi U, De Britto-Pereira MC. Frugivory and seed dispersal by *Hyla truncata*, a Neotropical treefrog. *Copeia*. 1989;1989:781. <https://doi.org/10.2307/1445517>.
105. De-Oliveira-Nogueira CH, Souza UF, Machado TM, Figueiredo-de-Andrade CA, Mônico AT, Sazima I, Sazima M, Toledo LF. Between fruits, flowers and nectar: the extraordinary diet of the frog *Xenohyla truncata*. *Food Webs*. 2023;35. <https://doi.org/10.1016/j.fooweb.2023.e00281>.
106. Schriever TA, Williams DD. Influence of pond hydroperiod, size, and community richness on food-chain length. *Freshw Sci*. 2013;32:964–75. <https://doi.org/10.1899/13-008.1>.
107. Tewksbury JJ, Anderson JGT, Bakker JD, Billo TJ, Dunwiddie PW, Groom MJ, Hampton SE, Herman SG, Levey DJ, Machnicki NJ, Del Rio CM, Power ME, Rowell K, Salomon AK, Stacey L, Trombulak SC, Wheeler TA. Natural history's place. *Sci Soc BioScience*. 2014;64:300–10. <https://doi.org/10.1093/biosci/biu032>.
108. Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J. Biodiversity hotspots for conservation priorities. *Nature*. 2000;403:853–8. <https://doi.org/10.1038/35002501>.
109. Violle C, Thuiller W, Mouquet N, Munoz F, Kraft NJB, Cadotte MW, Livingstone SW, Moullot D. Functional rarity: the ecology of outliers. *Trends Ecol Evol*. 2017;32:356–67. <https://doi.org/10.1016/j.tree.2017.02.002>.
110. Pimm SL, Jenkins CN, Abell R, Brooks TM, Gittleman JL, Joppa LN, Raven PH, Roberts CM, Sexton JO. The biodiversity of species and their rates of extinction, distribution, and protection. *Science*. 2014;344:1246752. <https://doi.org/10.1126/science.1246752>.
111. Cadotte MW, Carscadden K, Mirotnick N. Beyond species: functional diversity and the maintenance of ecological processes and services. *J Appl Ecol*. 2011;48:1079–87. <https://doi.org/10.1111/j.1365-2664.2011.02048.x>.
112. Noss RF. The naturalists are dying off. *Conserv Biol*. 1996;10:1–3.
113. Huckembeck S, Loebmann D, Albertoni EF, Hefler SM, Oliveira MCLM, Garcia AM. Feeding ecology and basal food sources that sustain the paradoxical frog *Pseudis minuta*: a multiple approach combining stomach content, prey availability, and stable isotopes. *Hydrobiologia*. 2014;740:253–64. <https://doi.org/10.1007/s10750-014-2022-2>.
114. Huckembeck S, Winemiller KO, Loebmann D, Garcia AM. Trophic ecology of two sympatric frogs with contrasting morphology and habitat use in a subtropical wetland. *Herpetologica*. 2018;74:207–16. <https://doi.org/10.1655/Herpetologica-a-D-17-00069.1>.
115. Lima AP, Moreira G. Effects of prey size and foraging mode on the ontogenetic change in feeding niche of *Colostethus stepheni* (Anura: Dendrobatidae). *Oecologia*. 1993;95:93–102. <https://doi.org/10.1007/BF00649512>.

116. Solé M, Beckmann O, Pelz B, Kwet A, Engels W. Stomach-flushing for diet analysis in anurans: an improved protocol evaluated in a case study in araucaria forests, Southern Brazil. *Stud Neotrop Fauna Environ.* 2005;40:23–8. <https://doi.org/10.1080/01650520400025704>.
117. Oliveira SSD, Silva ATD, Zina J. Diet of a bromeligenous species of treefrog in the Brazilian semiarid. *Braz J Biol.* 2021;81:411–7. <https://doi.org/10.1590/1519-6984.229742>.
118. Lima JE, de P, Rödder D, Solé M. Diet of two sympatric *Phyllomedusa* (Anura: Hylidae) species from a Cacao plantation in Southern bahia, Brazil. *North-West J Zool.* 2010;6:13–24.
119. Brandão RA, Fenker J, Lopes BEPDC, De Sena VMDA, Vasconcelos BD. Diet of terrestrial anurans in an ephemeral and simplified habitat during the dry season in the Brazilian Cerrado. *Ethol Ecol Evol.* 2020;32:527–50. <https://doi.org/10.1080/03949370.2020.1755373>.
120. Huning D, Tonial F, Oliveira, Zanella N, Brandalise J, de Guerra M, Ficanha K, Tedesco N CD. How does the feeding behaviour of the common forest Toad *Rhinella henseli* (Anura: Bufonidae) vary in space and time? Trophic ecology, chemical and antimicrobial activity. *Herpetol J.* 2022;32:59–69. <https://doi.org/10.33256/32.2.5969>.
121. Díaz JA, Sugai JLMM, Ceron K, Moroti M, de Santana T DJ. Dietary selectivity and sexual size dimorphism of *Chiasmocleis Mehelyi* (Anura: Microhylidae) in a Cerrado area of Southwest Brazil. *North-West J Zool.* 2020;16:166–71.
122. Meurer W, Gonçalves FG, Bovendorp RS, Percequillo AR, Bertoluci J. Diet electivity and preferences for food resources in *Chiasmocleis leucosticta* (Anura: Microhylidae). *J Herpetol.* 2021;55. <https://doi.org/10.1670/20-106>.
123. Traugott M, Kamenova S, Ruess L, Seeber J, Plantegenest M. Empirically characterising trophic networks. In: Woodward G, Bohan DA, editors. *Advances in ecological research.* Elsevier; 2013. pp. 177–224. <https://doi.org/10.1016/B978-0-12-420002-9.00003-2>.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.