

Predation of a Barker Frog, *Physalaemus cuvieri* Fitzinger, 1826, by a Giant Water Bug, genus *Belostoma*, in a Cerrado pond in Minas Gerais, Brazil

Wendell Sousa-Felix¹, Rayça Azevedo¹, Camila R.O. Leal¹, Priscila S. Carvalho², Diego J. Santana², and Karoline Ceron^{1,*}

Amphibians play a crucial role in ecosystems, due to their biphasic life cycle. They connect aquatic and terrestrial communities, where they serve as both predators and prey (Ceron et al., 2024). In aquatic environments, amphibian larvae are often herbivorous or planktivorous, and they help regulate the abundance and biomass of aquatic flora, including phytoplankton (West, 2018). Tadpoles have a significant impact as bioturbators, organisms that modify sediment distribution, stimulate microbial activity and nutrient absorption, and promote trophic dynamics and energy flow in an environment (Montaña et al., 2019). Adult frogs are also extremely important to the ecosystem, because they help regulate agricultural pest populations and human disease vectors (e.g., Peltzer and Lajmanovich, 2002; Valencia-Aguilar et al., 2013; Ceron et al., 2024). However, despite being predators in both aquatic and terrestrial environments, frogs are also prey for various groups of organisms, including carnivorous plants, vertebrates, and invertebrates (Duellman and Trueb, 1994; Toledo, 2007). Among invertebrates, frogs are known as prey for spiders (Barbo et al., 2009), ants (Kaczmarek et al., 2019), and water bugs (Hernández-Ruz et al., 2022), as well as other organisms.

Physalaemus cuvieri is a leptodactylid frog widely distributed in South America, with records from northern Uruguay, Argentina, eastern and northern Paraguay, Bolivia, Venezuela, and northeastern, central, and southern Brazil (Frost, 2025). This species can

be found in lentic and/or temporary waterbodies, and females are larger than males (Da Silva and Rossa-Feres, 2010). This widely distributed species has been reported as prey for diverse groups of animals, including spiders (e.g., genus *Ancylometes*; Maffei et al., 2010), water bugs (e.g., *Belostoma elongatum*; Toledo, 2003), snakes (e.g., *Erythrolamprus poecilogyrus*; Andrade et al., 2020), and another frogs (e.g., *Aquarana catesbeiana*; Boelter et al., 2012).

During a field expedition on 27 November 2019 to a Cerrado domain at Parque Nacional da Serra da Canastra, São Roque de Minas Municipality, Minas Gerais State, Brazil (20.2422°S, 46.5116°W; elevation 1386 m), we observed a giant water bug preying on a *P. cuvieri* in a perennial pond (Fig. 1A). Both specimens were collected and preserved in 70% ethanol (Fig. 1B). The frog was deposited in the Zoological Collection of the Mato Grosso do Sul Federal University in Campo Grande, Mato Grosso do Sul, Brazil (ZUFMS-AMP1743). The frog had a body length of 35.3 mm and the giant water bug, identified as a member of the genus *Belostoma* (Ribeiro, 1999) measured 35.3 mm. *Physalaemus cuvieri* can be identified by its dorsal coloration, which ranges from brownish to gray and features alternating light and dark stripes. A distinctive round light spot is present between the shoulders, along with a thick, dark band extending from the snout to the forelimbs; this band gradually narrows toward the inguinal region. The species also exhibits reddish inguinal and axillary regions. Males have a black gular region, while the venter is cream-colored or dark anteriorly, with cream vermiform markings (Uetanabaro et al., 2008). *Belostoma* was diagnosed by the following characters: the first rostral segment significantly longer than its greatest width and subequal in length to the second segment; a two-segmented foretarsus; and a well-developed hemelytral membrane with uniformly elongated narrow rectangular cells (Ribeiro, 1999; Estévez and Ribeiro, 2011).

¹ Laboratório de Interações Ecológicas e Biodiversidade, Departamento de Biologia, Universidade Federal do Ceará, Campus do Pici, Avenida Mister Hull s/n, Fortaleza, Ceará 60455-760, Brazil.

² Manguari - Laboratório de Biogeografia e Sistemática de Anfíbios e Répteis, Universidade Federal de Mato Grosso do Sul, Campo Grande, Mato Grosso do Sul 79070-900, Brazil.

* Corresponding author. E-mail: adenomera@gmail.com




Figure 1. (A) A giant water bug (genus *Belostoma*) preying on a Barker Frog (*Physalaemus cuvieri*) at Parque Nacional da Serra da Canastra, São Roque de Minas, Minas Gerais, Brazil. (B) Collected specimens and their relative sizes. Photos by Karoline Ceron (A) and Priscila S. Carvalho (B).

Our present observation reaffirms the potential of water bugs to prey on adult anurans, especially on males during their calling activity. In South America, the genus *Belostoma* comprises approximately 74 species, 40 of them recorded in Brazil (Ribeiro, 2007), in the states of Mato Grosso, Minas Gerais, Goiás, Santa Catarina, São Paulo, Rio de Janeiro, and Rio Grande do Sul (Figueiredo-de-Andrade et al., 2010). These aquatic insects regulate prey populations and serve as food source for various predators, also acting as bioindicators of environmental quality (Oliveira, 2019; Valbon et al., 2019; Hernández-Ruz et al., 2022). *Belostoma* species can act as natural biological control agents for disease vectors, such as the mosquitoes *Anopheles gambiae* (malaria vector), *Aedes aegypti* (dengue vector), and *Culex quinquefasciatus* (filariasis vector). These cosmopolitan bugs prey on both vertebrates and invertebrates (Valbon et al., 2019), often capturing organisms larger than themselves (Armúa and Estévez, 2006).

Giant water bugs are widely recognized as significant predators of anurans, preying on both larval and adult stages (Tobler and Plath, 2007). Studies have documented their predation on adult anurans, including adult treefrogs (genera *Dendropsophus* and *Scinax*), and juveniles of *Dendropsophus branneri*, *Leptodactylus pustulatus*, *L. pentadactylus*, and *L. labyrinthicus* (Toledo, 2003;

Baracho et al., 2014; Fadel et al., 2019; Hernández-Ruz et al., 2022). Despite the well-established predation pressure exerted by giant water bugs on tadpoles, little is known about whether the risk of predation from these insects shapes behavioural and morphological adaptations in adult anurans in aquatic environments, and the cues used by anurans to avoid predation. Assessing both consumptive and non-consumptive effects enhances our understanding of how giant water bugs shape anuran populations.

Acknowledgements. Specimens were collected under a permit from SISBIO (#66593-4). We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for financial support under Finance Code 001 (CROL). DJS thanks the Conselho Nacional de Desenvolvimento Científico e Tecnológico for his research fellowship (CNPq 311284/2023-0). KC supported by a research fellowship (#444240/2024-1) from the Conselho Nacional de Desenvolvimento Científico e Tecnológico.

References

- Andrade, H., da Costa, S.M., dos Santos, M.A., dos Reis Dias, E.J. (2020): Diet review of *Erythrolamprus poecilogyrus* (Wied-Neuwied, 1825) (Serpentes: Dipsadidae), and first record of *Dermatonotus muelleri* (Boettger, 1885) (Anura: Microhylidae) as a prey item in Sergipe State, northeastern Brazil. *Herpetology Notes* **13**: 1065–1068.
- Armúa de Reyes, C.A., Estévez, A.L. (2006): Predation on *Biomphalaria* sp. (Mollusca: Planorbidae) by three species of

- the genus *Belostoma* (Heteroptera: Belostomatidae). Brazilian Journal of Biology **66**: 1033–1035.
- Baracho, E.B.O., Silva, J.S., Nascimento, B.H.M., Fonseca, E.M.F., Magalhães, F.M. (2014): *Dendropsophus branneri* (Cochran, 1948) (Anura: Hylidae) as prey to invertebrates in northeastern Brazil. Herpetology Notes **7**: 17–19.
- Barbo, F.E., Rodrigues, M.G., Couto, F.M., Sawaya, R.J. (2009): Predation on *Leptodactylus marmoratus* (Anura: Leptodactylidae) by the spider *Ctenus medius* (Araneae: Ctenidae) in the Atlantic Forest, southeast Brazil. Herpetology Notes **2**: 99–100.
- Boelter, R.A., Kaefer, I.L., Both, C., Cechin, S. (2012): Invasive bullfrogs as predators in a Neotropical assemblage: what frog species do they eat? Animal Biology **62**: 397–408.
- Ceron, K., Santana, S.J., Pires, M.M. (2024): The economic risk of the losses in pest control as frogs decline. EcoEvoRxiv. Available at: <https://ecoevorxiv.org/repository/view/7363>. Accessed on 29 August 2025.
- da Silva, F.R., Rossa-Feres, D.D.C. (2010): Seasonal variation in body size of tropical anuran amphibians. Herpetology Notes **3**: 205–209.
- Duellman, W.E., Trueb, L. (1994): Biology of Amphibians. Baltimore, Maryland, USA, The Johns Hopkins University Press.
- Estévez, A.L., Ribeiro, J.R.I. (2011): *Weberiella* De Carlo, 1966 (Insecta: Heteroptera: Belostomatidae) revisited: redescription with a key to the genera of Belostomatidae and considerations on back-brooding behaviour. Zoologischer Anzeiger **250**: 46–54.
- Fadel, R.M., Thaler, R., Folly, H., Galvão, C., Hoffmann, M., da Silva, L.A., et al. (2019): Predation of anurans across multiple life stages in an Amazon-Cerrado transitional zone. Herpetology Notes **12**: 895–899.
- Figueiredo-de-Andrade, C.A., Santana, D.J., de Carvalho-e-Silva, S.P. (2010): Predation on *Scinax x-signatus* (Anura: Hylidae) by the giant water bug *Lethocerus annulipes* (Hemiptera: Belostomatidae) in a Brazilian Restinga habitat. Amphibia-Reptilia **14**: 307–309.
- Frost, D.R. (2025): Amphibian Species of the World: an Online Reference. Version 6.2. Available at: <https://amphibiansoftheworld.amnh.org/index.php>. Accessed on 29 August 2025.
- Hernández-Ruz, E.J., do Nascimento, F.O.P., da Silva, G.A., da Silva, W.G., Correa, J.K.C. (2022): Predation on anurans by a giant water bug *Belostoma* sp. (Hemiptera: Belostomatidae) in a forest management area in the eastern Amazon. Food Webs **32**: e00238.
- Kaczmarek, M., Michlewicz, M., Tryjanowski, P. (2019): Ants biting amphibians: a review and new observations. European Journal of Ecology **5**: 118–125.
- Maffei, F., Ubaid, F.K., Jim, J. (2010): Predation of herps by spiders (Araneae) in the Brazilian Cerrado. Herpetology Notes **3**: 167–170.
- Montaña, C.G., Silva, S.D., Hagyar, D., Wager, J., Tieg, L., Sadeghian, C., et al. (2019): Revisiting “what do tadpoles really eat?” A 10-year perspective. Freshwater Biology **64**: 2269–2282.
- Oliveira, N.R.X.D. (2019): Alterações comportamentais e fisiológicas em baratas d'água, *Belostoma anurum* (Hemiptera: Belostomatidae) decorrentes da exposição ao cádmio. Unpublished MSc thesis, Universidade Federal de Viçosa, Viçosa, Brazil.
- Peltzer, P.M., Lajmanovich, R.C. (2002): Preliminary studies of food habits of *Lysapsus limellus* (Anura, Pseudidae) in lentic environments of Parana River, Argentina. Bulletin de la Société Herpétologique de France **101**: 53–58.
- Ribeiro, J.R.I. (1999): Revisão das Espécies de *Belostoma* Latreille, 1807 (Insecta: Heteroptera: Belostomatidae) ocorrentes no Sudeste do Brasil. Unpublished MSc thesis, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.
- Ribeiro, J.R.I. (2007): Uma revisão das espécies de *Belostoma* Latreille, 1807 (Hemiptera: Heteroptera: Belostomatidae) dos quatro estados do sudeste brasileiro. Zootaxa **1807**: 1–70.
- Tobler, M., Schlupp, I., Plath, M. (2007): Predation of a cave fish (*Poecilia mexicana*, Poeciliidae) by a giant water-bug (*Belostoma*, Belostomatidae) in a Mexican sulphur cave. Ecological Entomology **32**: 492–495.
- Toledo, L.F. (2003): Predation on seven South American anuran species by water bugs (Belostomatidae). Phyllomedusa **2**: 105–108.
- Toledo, L.F. (2007): Predação e defesa de anuros: revisão, descrição e evolução. Unpublished PhD thesis, Universidade Estadual de São Paulo, São Paulo, Brazil.
- Uetanabaro, M., Prado, C.P.A., Rodrigues, D.D.J., Gordo, M., Campos, Z. (2008): Guia de Campo de Anuros do Pantanal Sul e Planaltos de Entorno. Campo Grande, Mato Grosso do Sul, Brazil, Editora UFMS/UFMT.
- Valbon, W.R., Haddi, K., Gutiérrez, Y., Cruz, F.M., Azevedo, K.E.X., Perez, J.S.C., et al. (2019): Life history traits and predatory performance of *Belostoma anurum* (Hemiptera: Belostomatidae), a biological control agent of disease vector mosquitoes. Neotropical Entomology **48**: 899–908.
- Valencia-Aguilar, A., Cortés-Gómez, A.M., Ruiz-Agudelo, C.A. (2013): Ecosystem services provided by amphibians and reptiles in neotropical ecosystems. International Journal of Biodiversity Science, Ecosystem Services and Management **9**: 257–272.
- West, J. (2018): Importance of amphibians: a synthesis of their environmental functions, benefits to humans, and need for conservation. Unpublished Honors Thesis, Bridgewater State University, Bridgewater, Massachusetts, USA. Available at: https://vc.bridgew.edu/honors_proj/261. Accessed on 29 August 2025.