

NOTES ON GEOGRAPHIC DISTRIBUTION

**Amphibia, Anura, Hylidae, *Scinax auratus*:  
Distribution extension, new state records, and distribution map**

Diego José Santana<sup>1</sup>  
Henrique Caldeira Costa<sup>1</sup>  
Leandro de Oliveira Drummond<sup>3</sup>  
Paula Leão Ferreira<sup>2</sup>  
Renato Neves Feio<sup>1</sup>

<sup>1</sup> Universidade Federal de Viçosa, Museu de Zoologia João Moojen.  
Vila Giannetti 32. CEP 36570-000, Viçosa, MG, Brazil. E-mail: santana\_herpeto@yahoo.com.br.

<sup>2</sup> Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis.  
Parque Nacional Grande Sertão Veredas. CEP 39314-000. Chapada Gaúcha, MG, Brazil.

<sup>3</sup> Universidade Federal de Ouro Preto, Departamento de Ciências Biológicas, Laboratório de Zoologia dos Vertebrados. Campus Morro do Cruzeiro, CEP 35400-000. Ouro Preto, MG, Brazil

*Scinax auratus* (Wied-Neuwied, 1821) (Figures 1 and 2) is a small hylid treefrog associated with the *Scinax ruber* clade (*sensu* Faivovich 2002), occurring in the Atlantic Forest of northeastern Brazil (Alves et al. 2004; Frost 2009). This species inhabits forests and rocky outcrops along the border of forested and open areas (Alves et al. 2004). It is generally found inside terrestrial and epiphytic bromeliads, or calling inside bushes and grassy vegetation associated to lentic water bodies, where it reproduces (Bokermann 1969).


Brazil (Bastazini et al. 2007; Bokermann 1969; Nunes et al. 2007; Juncá 2006), and Quebrangulo, state of Alagoas (Alves et al. 2004).

In the present paper we report new records for *Scinax auratus* and a map representing the distribution of this species (Figure 3) based on published data, specimens collected by the authors, and specimens deposited in the following Brazilian herpetological collections: *Museu de Zoologia João Moojen, Universidade Federal de Viçosa, Viçosa, state of Minas Gerais (MZUFV); Laboratório de Zoologia dos Vertebrados, Universidade Federal de Ouro Preto, Ouro Preto, state of Minas Gerais (LZV/UFOP); and Museu de Zoologia da Universidade Federal da Bahia, Salvador, state of Bahia (UFBA).*




**Figure 1.** *Scinax auratus* (MZUFV 5473, male) from *Fazenda Santana*, municipality of Salto da Divisa, Minas Gerais, Brazil. Photo by R. N. Feio.

Literature data reported the occurrence of this species in the localities of Camaçari, Feira de Santana, Mata de São João, Maracás (type locality), and Santa Inês, all in the state of Bahia,



**Figure 2.** *Scinax auratus* (LZV-A 1146, male) from *Reserva Biológica de Santa Isabel*, municipality of Pirambú, Sergipe, Brazil. Photo by L. O Drummond.

### NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 3.** Geographic distribution map of *Scinax auratus*. ALAGOAS (AL): 1 - Quebrangulo; SERGIPE (SE): 2 - Pirambú; BAHIA (BA): 3 - Entre Rios; 4 - Feira de Santana; 5 - Mata de São João; 6 - Camaçari; 7 - Cachoeira; 8 - Salvador; 9 - Amargosa; 10 - Santa Inês; 11 - Maracás; MINAS GERAIS (MG): 12 - Salto da Divisa.

We identified several new localities of occurrence of *Scinax auratus*, including two new state records. Specimens from the following municipalities in the state of Bahia, Brazil, represent new localities for the species: Amargosa (13°01' S, 39°36' W; UFBA 6481, 6487), Cachoeira (12°44' S, 38°52' W; UFBA 7582-7583), Entre Rios (11°57' S, 38°04' W; UFBA 5970, 6138), and Salvador (12°58' S, 38°30' W; UFBA 432-436, UFBA 438-447, UFBA 6060).

During a fieldwork in Reserva Biológica de Santa Isabel, municipality of Pirambú (10°44' S, 36°51' W), northern coast of the state of Sergipe, several individuals of *Scinax auratus* (vouchers LZV-A 1146-1147, 1194-1197) were observed in areas of "restinga", a phytophysionomy of the Atlantic Forest biome typical of the sandy coastal plains that occupies 79% of the Brazilian coast (Araujo et al. 1998). Males were found calling on the edge or on the emergent vegetation of ponds surrounded by arboreal and anthropized *restingas*.

This is the first record of this species for the state of Sergipe, filling a distributional gap between the states of Bahia and Alagoas.

The occurrence of *S. auratus* at *Fazenda Santana*, municipality of Salto da Divisa, state of Minas Gerais (16°00' S, 39°56' W; MZUFV 4908-4910, 5472-5473) provides another new state record for the species and its southernmost record, ca. 290 km south from the municipality of Maracás (type locality), state of Bahia (Bokermann 1969).

The current data on the geographic distribution of *S. auratus* shows that this species occurs along the Atlantic Forest domain (sensu Ab'Sáber 1977), from Quebrangulo, in the state of Alagoas, to Salto da Divisa, in the northeastern region of the state of Minas Gerais (Figure 3). The distributional gap between Maracás (BA) and Salto da Divisa (MG) is probably due to insufficient sampling.

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Acknowledgements

We thank Dr. Marcelo F. Napoli (UFBA) for allowing the examination of specimens deposited in UFBA and critical reading on a previous version of the manuscript. Luciana B. Nascimento (*Museu de Ciências Naturais, Pontifícia Universidade Católica de Minas Gerais*, Belo Horizonte, Minas Gerais) and Mirco Solé (*Universidade Estadual de Santa Cruz*, Ilhéus, Bahia) also provided insightful comments on this work. Mário R. Moura kindly made the map. The specimens from Minas Gerais were collected during the project “*Inventário de áreas prioritárias para conservação da biodiversidade nos vales dos rios Jequitinhonha e Mucuri*”, supported by the Project on the Conservation and Sustainable Use of Brazilian Biological Diversity/Brazilian Ministry of the Environment (PROBIO/MMA) and coordinated by Conservation International do Brasil (CI-Brasil) (collection permit # 199/2002/FAUNA-MG; process # 02015.018836/02). The specimens from Sergipe were collected during the elaboration of the Management Plan of *Reserva Biológica de Santa Isabel*, coordinated by *Instituto Chico Mendes de Conservação da Biodiversidade* (ICMBio) and supported by *Fundação do Amparo à Pesquisa do Estado de Minas Gerais* (FAPEMIG) (collection permit number 13089-1).

---

### Literature cited

- Ab'Sáber, A. N. 1977. Os domínios morfoclimáticos na América do Sul. Primeira aproximação. *Geomorfologia* 52:1-21.
- Alves, A. C. R., M. R. Gomes, and S. P. Carvalho-e-Silva. 2004. Description of the tadpole of *Scinax auratus* (Wied-Neuwied) (Anura, Hylidae). *Revista Brasileira de Zoologia* 21(2):315-317.
- Araujo, D. S. D., F. R. Scarano, C. F. C. Sá, B. C. Kurtz, H. L. T. Zaluar, R. C. M. Montezuma, and R. C. Oliveira. 1998. Comunidades vegetais do Parque Nacional da Restinga de Jurubatiba; p. 39-62 *In* F. A. Esteves (ed.). *Ecologia das Lagoas costeiras do Parque Nacional da Restinga de Jurubatiba e do Município de Macaé* (RJ). Rio de Janeiro: UFRJ.
- Bastazini, C. V., J. V. Munduruca, P. L. B. Rocha, and M. F. Napoli. 2007. Which environmental variables better explain changes in anuran community composition? A case study in the restinga of mata de São João, Bahia, Brazil. *Herpetologica* 63(4): 459-471.
- Bokermann, W. C. A. 1969. Notas sobre *Hyla aurata* Wied, 1824 (Anura, Hylidae). *Revista Brasileira de Biologia* 29(2):159-162.
- Faivovich, J. 2002. A cladistic analysis os *Scinax* (Anura: Hylidae). *Cladistics* 18(2002): 367-393.
- Frost, D. 2009. Amphibian Species of the World: an Online Reference. Version 5.3. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/> American Museum of Natural History, New York, USA. Captured on April 2009.
- Juncá, F. A. 2006. Diversidade e uso de hábitat por anfíbios anuros em duas localidades de Mata Atlântica, no norte do estado da Bahia. *Biota Neotropica* 6(2): 1-17.
- Nunes, I., R. S. Santiago, and F. A. Juncá. 2007. Advertisement calls of four hylid frogs from the State of Bahia, northeastern Brazil (Amphibia, Anura, Hylidae). *South American Journal of Herpetology* 2(2): 89-96.

Received November 2008

Accepted May 2009

Published online May 2009