

*Thamnodynastes hypoconia*  
(COPE, 1860), preys upon *Scinax*  
*fuscomarginatus* (LUTZ, 1925)

Knowledge on feeding habits contributes to the understanding of a species' natural history as well as interactions and energy flux through ecosystems (FALICO et al. 2012; LINARES et al. 2016). Amphibians play important roles in food chains of ecosystems in representing a link between terrestrial and aquatic environments (MACHADO et al. 2015). However, predation events on amphibians are only rarely observed in nature (POMBAL 2007).

The frog genus *Scinax* WAGLER, 1830, is composed of 71 species, 51 of which occur in Brazil (FROST 2017). *Scinax fuscomarginatus* (LUTZ, 1925) inhabits open areas of the Pantanal and the Cerrado ecoregions (BRASILEIRO et al. 2005). Despite being widely distributed, records of predation upon it are scarce, as is the knowledge of its predators (BRANDÃO & GARDA 2000; ROLIM et al. 2009).

*Thamnodynastes hypoconia* (COPE, 1860) is a small snake with semi-arboreal habits, frequently reported in flooded envi-



Fig. 1. A - Young *Thamnodynastes hypoconia* (COPE, 1860) (ZUFMS 2477; snout-vent-length 191 mm) devouring a small adult *Scinax fuscomarginatus* (LUTZ, 1925) (ZUFMS 5626; SVL 22.9 mm); B - Stomach opened to identify the ingested anuran species.

ronments (SAWAYA et al. 2008; ARAUJO et al. 2010). Its diet includes anurans, fishes and lizards (MARQUES et al. 2001; SAWAYA et al. 2008; ARAUJO et al. 2010; MAFFEI et al. 2011; MOYA & MAFFEI 2012). It is a predator to several species of hyloid frogs (see

DORIGO et al. 2014). The present note is the first that reports a predation event by *T. hypoconia* upon an individual of *S. fuscomarginatus*.

On May 15, 2017, at 20:40 h, during herpetological field work in the Reserva

Particular do Patrimônio Natural (RPPN) Fazenda Santa Fé, municipality of Campo Grande, Mato Grosso do Sul (20.5131 S, 54.7277 W, 500 m a.s.l), the authors observed a young *T. hypoconia* (snout-vent-length 191 mm) preying upon an adult *S. fuscomarginatus* (SVL 22.9 mm). The snake was detected in a swamp area, sitting on emergent herbaceous vegetation, about 50 cm above the water surface. The capture proper of the prey was not observed, the snake was about to swallow the treefrog (Fig. 1A). The end of the ingestion was waited for, the snake was caught, euthanased and analyzed in the laboratory for its stomach content (Fig. 1B). Both specimens collected (Collection Permit SISBIO 45889-1) are housed in the Coleção Zoológica da Universidade Federal de Mato Grosso do Sul (*T. hypoconia* ZUFMS-REP 2477; *S. fuscomarginatus* ZUFMS-AMP 5626).

Anurans are commonly preyed on by snakes because they are abundant in most of the habitats and the snake's risk of injuries by retaliation is low (MARTINS et al. 2002; TOLEDO et al. 2007). Besides, this predation event occurred at a typical calling site of *S. fuscomarginatus* (see TOLEDO & HADDAD 2005), where and when males are probably particularly exposed to predation.

**ACKNOWLEDGMENTS:** The authors are grateful to Marina and João for giving their permission to study the area of RPPN Fazenda Santa Fé; M. de Toledo Moroti, K. Ceron and J. F. Cuestas Carrillo are grateful to CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for scholarships.

**REFERENCES:** ARAUJO, C. O. & CORRÊA-FILHO, D. T. & SAWAYA, R. J. (2010): Serpentes da estação ecológica de Santa Bárbara, SP: Um remanescente de cerrado do sudeste do Brasil.- *Biota Neotropica*, São Paulo; 10: 235-245. BRANDÃO, R. A. & GARDA, A. A. (2000): *Scinax fuscomarginatus* (NCN). Predation.- *Herpetological Review*, New York; 31 (3):171. BRASILEIRO, C. A. & SAWAYA, R. J. & KIEFFER, M. C. & MARTINS, M. (2005): Amphibians of an open Cerrado fragment in southeastern Brazil.- *Biota Neotropica*, São Paulo; 5 (2): 93-109. DORIGO, T. A. & VRCIBRADIC, D. & BORGES-JUNIOR, V. N. T. (2014): New records of anuran predation by snakes of the genus *Thamnodynastes* WAGLER, 1830 (Colubridae: Dipsadinae) in the Atlantic rainforest of southeastern Brazil.- *Herpetology Notes*, Braunschweig; 7: 261-264. FALICO, D. A. & LÓPEZ, J. A. & ANTONIAZZI, C. E. (2012): Opportunistic predation upon dragonflies by *Pseudis limellum* and *Pseudis paradoxa* (Anura: Hylidae) in the Gran Chaco region, Argentina.- *Herpetology Notes*, Braunschweig; 5: 215-217. FROST, D. R. (2017): Amphibian Species of the World: an Online Reference. Version 6.0. WWW resource

available at < <http://research.amnh.org/herpetology/amphibia/index.html> >. [last accessed on May 2, 2017], American Museum of Natural History, New York. LINARES, A. M. & MACIEL-JUNIOR, J. A. H. & MELLO, H. E. S. & LEITE, F. S. F. (2016): First report on predation of adult anurans by Odonata larvae.- *Salamandra*, Mannheim; 52 (1): 42-44. MACHADO, I. F. & MENEGUCCI, R. C. & MENDES, H. F. & MOROTI, M. T. (2015): Polyphenism: Defensive colour behaviour of *Phasmahyla guttata* (A. LUTZ, 1924) (Amphibia, Anura, Hylidae).- *Herpetology Notes*, Braunschweig; 8: 467-470. MAFFEL, F. & ROLIM, D. C. & JIM, J. (2011): *Thamnodynastes hypoconia*. Diet.- *Herpetological Review*, New York; 42: 620. MARQUES, O. A. V. & ETEROVIC, A. & SAZIMA, I. (2001): Serpentes da Mata Atlântica: guia ilustrado para a Serra do Mar. Ribeirão Preto, Brazil (Holos), pp.184. MARTINS, M. & MARQUES, O. A. V. & SAZIMA, I. (2002): Ecological and phylogenetic correlates of feeding habits in Neotropical pitvipers of the genus *Bothrops*; pp. 307-328. In: SCHUETT, G. W. & HOGGREN, M. & DOUGLAS, M. E. & GREENE, H. W. (Eds.): *Biology of the Vipers*. Eagle Mountain (Eagle Mountain Publ.). MOYA, G. M. & MAFFEL, F. (2012): Predation on *Dendropsophus elianeae* (NAPOLI & CARAMASCHI, 2000) (Anura: Hylidae) by *Thamnodynastes hypoconia* (COPE, 1860) (Squamata: Colubridae).- *Herpetology Notes*, Braunschweig; 5: 261-262. POMBAL J. P. (2007): Notas sobre predação em uma taxocenose de anfíbios anuros no sudeste do Brasil.- *Revista Brasileira de Zoologia*, Curitiba; 24: 841-843. ROLIM, D. C. & DE ALMEIDA, S. C. & JIM, J. (2009): *Scinax fuscomarginatus* (NCN). Predation.- *Herpetological Review*, New York; 40 (1): 73. SAWAYA, R. J. & MARQUES, O. A. V. & MARTINS, M. (2008): Composition and natural history of a Cerrado snake assemblage at Itirapina, São Paulo State, southeastern Brazil.- *Biota Neotropica*, São Paulo; 8: 127-149. TOLEDO, L. F. & HADDAD, C. F. B. (2005): Acoustic repertoire and calling behavior of *Scinax fuscomarginatus* (Anura, Hylidae).- *Journal of Herpetology*, Houston, etc.; 39: 455-464. TOLEDO, L. F. & RIBEIRO, R. S. & HADDAD, C. F. B. (2007): Anurans as prey: an exploratory analysis and size relationships between predators and their prey.- *Journal of Zoology*, London, etc.; 271: 170-177.

**KEY WORDS:** Reptilia: Squamata: Serpentes: Colubridae; *Thamnodynastes hypoconia*. Amphibia: Anura: Hylidae; *Scinax fuscomarginatus*, predation, food ecology, biology; Mato Grosso do Sul, Brazil, South America

**SUBMITTED:** July 31, 2017

**AUTHORS:** João Lucas Lago CANHETE <sup>1)</sup>, Matheus DE TOLEDO MOROTI <sup>1)</sup>, Juan Fernando CUESTAS CARRILLO <sup>2)</sup>, Karoline CERON <sup>2)</sup> & Diego José SANTANA (Corresponding author < [santanadiegojose@yahoo.com](mailto:santanadiegojose@yahoo.com) >). <sup>1, 2)</sup>

<sup>1)</sup> Matinguari – Laboratório de Sistemática e Biogeografia de anfíbios e répteis, Instituto de Biociências, Universidade Federal de Mato Grosso do Sul, Cidade Universitária, CEP 79002-970, Campo Grande, Mato Grosso do Sul, Brazil.

<sup>2)</sup> Programa de Pós-Graduação em Ecologia e Conservação, Instituto de Biociências, Universidade Federal de Mato Grosso do Sul, Cidade Universitária, CEP 79002-970, Campo Grande, Mato Grosso do Sul, Brazil.