

HELICOPS ANGULATUS (Watersnake). DIET. *Helicops angulatus* occurs in northern South America, with records from Atlantic forest or Amazonian environments in Colombia, Venezuela, Trinidad, Guiana, Suriname, French Guiana, Ecuador, Peru, Bolivia, and northern and northeastern regions of Brazil (França et al. 2006. Occ. Pap. Sam Noble Oklahoma Mus. Nat. Hist. 17:1–3; Infante-Rivero et al. 2008. Herpetotropicos 4:39–39). They are

aquatic, nocturnal, and feed on fish, tadpoles, and occasionally adult anurans (Ford et al. 2002. Carib. J. Sci. 38:129–132; Scartozzoni 2009. Unpubl. DS thesis. Universidade de São Paulo, São Paulo).

A mature female *H. angulatus* (SVL = 297 mm) was captured on 9 November 2010, at 2200 h, on the edge of a stream in Atlantic Forest of Cruz do Espírito Santo municipality, Paraíba State, northeastern Brazil. Dissection revealed remains of an adult *Leptodactylus natalensis* (SVL = 49.4 mm) in the snake's stomach. This is the first record of consumption of *L. natalensis* by *H. angulatus*. Because both species are sympatric and their habitats are similar, this type of interaction may be expected. The habit of making nests and vocalizing at the edge of slow-flowing streams (Amorim et al. 2009. Pap. Avul. Zool. 49:1–7) may make *L. natalensis* more vulnerable to predators such as *H. angulatus* during the breeding season. The snake and frog are housed in the Coleção Herpetológica da Universidade Federal da Paraíba, João Pessoa (CHUFPB 00005).

We thank the Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA) for permits. DJS would like to thank CAPES for their current Ph.D. scholarships, CAPES for scholarship to ASP and RR, and CNPq for scholarship to RLA and research fellowship to DOM. FGRF thanks the financial support from the CNPq (Universal grant 474250/2010-5). DOM thanks the financial support from the CNPq (Universal grant 481537/2009-0).

RALPH LACERDA DE ALBUQUERQUE (e-mail: ralph.la@gmail.com),
DANIEL ORSI LARANJEIRAS (e-mail: danorsi@gmail.com), **ARIELSON DOS SANTOS PROTÁZIO** (e-mail: neu_ptz@hotmail.com), **DIEGO JOSÉ SANTANA** (e-mail: santana_herpeto@yahoo.com.br), **RICARDO RODRIGUES** (e-mail: ricardodasilveira@gmail.com), **DANIEL OLIVEIRA MESQUITA** (e-mail: danmesq@dse.ufpb.br), Universidade Federal da Paraíba, Centro de Ciências Exatas e da Natureza, Departamento de Sistemática e Ecologia, Laboratório de Herpetologia, Cidade Universitária, CEP 58059-900, João Pessoa, Paraíba, PB, Brazil; **FREDERICO GUSTAVO RODRIGUES FRANÇA**, Universidade Federal da Paraíba, Centro de Ciências Aplicadas e Educação, Campus IV, Litoral Norte Rua da Mangueira, s/n, CEP 58297-000, Rio Tinto, PB, Brazil (e-mail: fredericogr@gmail.com).