

# Anurans from an Atlantic Forest-Caatinga ecotone in Rio Grande do Norte State, Brazil

Felipe de Medeiros Magalhães<sup>1</sup>, Anne Karenine Bezerra da Penha Dantas<sup>1</sup>, Marcos Roberto Monteiro de Brito<sup>1</sup>, Pedro Henrique Silveira de Medeiros<sup>1</sup>, Alan Felipe Oliveira<sup>1</sup>, Thiago César Sena de Oliveira Pereira<sup>1</sup>, Marcelo Henrique Campos de Queiroz<sup>1</sup>, Diego José Santana<sup>2</sup>, Willianilson Pessoa da Silva<sup>1</sup>, and Adrian Antonio Garda<sup>1,\*</sup>

**Abstract.** Brazil harbours the highest richness of anuran amphibians, and such diversity may be explained by the extensive and heterogeneous biomes and habitats occurring in this country. Still, the knowledge about anuran diversity and distribution in Brazil is marked by large gaps, and such data paucity hampers scientifically based decisions for species management and conservation, especially in north-eastern Brazil. This fact is clearly exemplified by Rio Grande do Norte State, where no species survey publication exists. Given this large gap, we surveyed an Atlantic Forest-Caatinga transition zone for thirteen months, in the municipality of Macaíba, Rio Grande do Norte State. We identified 34 amphibian anuran species at the locality, several of which are presented herein as the first published record for the State.

**Keywords.** Amphibians, Frogs, distribution, Escola Agrícola de Jundiaí, Macaíba.

## Introduction

The Neotropical region harbours the highest diversity of anuran amphibians, with about 2000 species or approximately one third of the world's known species (Duellman, 1999). The Brazilian territory hosts the highest richness of amphibians, with 946 known species (Segalla et al., 2012). This high diversity may be explained by the extensive landscape domains, each formed by collections of fairly heterogeneous and unique ecosystems in terms of their ecological features (Ab'Saber, 1977).

The Atlantic Forest in north-eastern Brazil comprises an eastern narrow zone of coastal forests adjacent to the Caatinga, a mosaic of xerophytic, deciduous, and semiarid thorn scrub vegetation (Ab'Saber, 1977). Most faunal diversity studies in the region have focused on sites under the influence of one of these biomes, while seasonal deciduous forests (characteristics of narrow transition zones between these two biomes), have received less attention.

Studies on transition zones are relevant because of

the ecological uniqueness, and hence conservation importance of these areas (Remanamanjato et al., 2002). Because transition zones may exhibit physical and biological characteristics of both adjacent regions (Willians, 1996), they may present a unique composition including species sets derived from both domains, and such sets can often be richer than either biome alone (Remanamanjato et al., 2002).

The knowledge about amphibian distributions, ranges, and community composition in north-eastern of Brazil is marked by large gaps and incomplete inventories (Rodrigues, 2003; Tabarelli and Silva, 2003). Such data paucity severely hampers conservation initiatives in highly diverse and threatened biome, such as the Atlantic Forest (Mittermeier et al., 1999). Basic data on distribution and community composition are crucial to define priority areas for biodiversity conservation, as well to manage and monitor species in the long-term (Camardelli and Napoli, 2012). This situation is so critical that not a single species list has been published for large areas in Brazil, such as Rio Grande do Norte State (with an area of 52.000 km<sup>2</sup>, larger than the Netherlands, for example). Even with recent research conducted in the region (Garda et al., 2010; Magalhães et al., 2012; Santana et al., 2011; São-Pedro et al., 2011), the assessment on anuran community composition in specific sites remains incipient.

This information gap in Rio Grande do Norte (RN) State persists despite the old colonization of the region, the extensive fragmentation and habitat loss (only 10%

<sup>1</sup> Universidade Federal do Rio Grande do Norte, Departamento de Botânica, Ecologia e Zoologia, Laboratório de Anfíbios e Répteis-LAR, CEP 59078-970, Natal, RN, Brazil.

<sup>2</sup> Universidade Federal da Paraíba, Centro de Ciências Exatas e da Natureza, Departamento de Sistemática e Ecologia, CEP 58051-900, João Pessoa, PB, Brazil.

\* Corresponding author. E-mail: pseudis@gmail.com

**Table 1.** Amphibian species recorded in Escola Agrícola de Jundiaí (EAJ), Macaéba Municipality, Rio Grande do Norte State, Brazil.


Brachycephalidae	<i>Ischnocnema ramagii</i> (Boulenger, 1888)
Bufonidae	<i>Rhinella granulosa</i> (Spix, 1824)
	<i>Rhinella jimi</i> (Stevaux, 2002)
Cycloramphidae	<i>Proceratophrys cristiceps</i> (Müller, 1883)
Hylidae	<i>Corythomantis greeningi</i> Boulenger, 1896
	<i>Dendropsophus branneri</i> (Cochran, 1948)
	<i>Dendropsophus minutus</i> (Peters, 1872)
	<i>Dendropsophus nanus</i> (Boulenger, 1889)
	<i>Dendropsophus oliveirai</i> (Bokermann, 1963)
	<i>Dendropsophus soaresi</i> (Caramaschi & Jim, 1983)
	<i>Hypsiboas albomarginatus</i> (Spix, 1824)
	<i>Hypsiboas raniceps</i> (Cope, 1862)
	<i>Phyllomedusa nordestina</i> Caramaschi, 2006
	<i>Scinax cretetus</i> Nunes & Pombal, 2011
	<i>Scinax fuscomarginatus</i> (A. Lutz, 1925)
	<i>Scinax cf. nebulosus</i> (Spix, 1824)
	<i>Scinax pachychirus</i> (Miranda-Ribeiro, 1937)
	<i>Scinax x-signatus</i> (Spix, 1824)
Leptodactylidae	<i>Leptodactylus caatingae</i> Heyer & Juncá, 2003
	<i>Leptodactylus fuscus</i> (Schneider, 1799)
	<i>Leptodactylus macrosternum</i> Miranda-Ribeiro, 1926
	<i>Leptodactylus mystaceus</i> (Spix, 1824)
	<i>Leptodactylus natalensis</i> A. Lutz, 1930
	<i>Leptodactylus troglodytes</i> A. Lutz, 1926
	<i>Leptodactylus vastus</i> A. Lutz, 1930
	<i>Leptodactylus</i> sp. (gr. <i>marmoratus</i> )
	<i>Physalaemus albifrons</i> (Spix, 1824)
	<i>Physalaemus cuvieri</i> Fitzinger, 1826
	<i>Pleurodema diplolister</i> (Peters, 1870)
	<i>Pseudopaludicola cf. falcipes</i>
	<i>Pseudopaludicola</i> sp.
Microhylidae	<i>Elachistocleis cesarii</i> (Miranda-Ribeiro, 1920)
	<i>Dermatonotus muelleri</i> (Boettger, 1885)
Ranidae	<i>Lithobates palmipes</i> (Spix, 1824)

of the original cover remains) caused by pastures and sugar cane cultivation, and the intensive knowledge on amphibian communities in neighbouring states (Borges-Nojosa and Cascon, 2005; Borges-Nojosa et al., 2010; Moura, 2010; Santana, 2010; Santana et al., 2008). Fragmentation and habitat loss can lead to dramatic reductions in species diversity (Cushman, 2006), but the lack of information on species compositions precludes an evaluation of their impacts in RN. More importantly, the lack of information impedes scientifically based decisions for the local fauna management and conservation. Thus, as part of a larger effort to fill the information gap on

anurans in RN, we surveyed for thirteen months an anuran community within an Atlantic Forest—Caatinga transition area in RN. We discuss species distribution extensions and contextualize the results in face of the richness reported for other areas in north-eastern Brazil.

## Material and Methods

The study was conducted at Escola Agrícola de Jundiaí (EAJ), municipality of Macaéba, Rio Grande do Norte State ( $5^{\circ} 53' 06.68''S$   $35^{\circ} 22' 01.28''W$ , Figure 1). The local climate is transitional between As' and BSh' types, according to Köppen classification (Cestaro and Soares, 2004). Deciduous lowland



**Figure 1.** The study site, Escola Agrícola de Jundiaí (black dot), located in the transition zone of the Atlantic Forest and Caatinga domains in Rio Grande do Norte State, north-eastern Brazil.


forests with both Caatinga and Atlantic forest domain characterize the region. Several temporary ponds form in the area during the rainy season, which also contains a permanent lake and a shallow permanent river (Jundiaí River) with sandy bottom. Human activity is intense in the region with the presence of cattle and pastures surrounding the study area.

We conducted field expeditions from April 2009 to May 2010. Most of the field effort occurred during the raining season (March to July). Rainfall data were obtained from EMPARN (Empresa de Pesquisa Agropecuária do Rio Grande do Norte), and local average is around 1200 mm. We actively searched for adults and tadpoles following a 2 km transect that crossed several temporary and semi-permanent ponds. Voucher specimens, photographs, and advertisement calls for at least one individual per species were collected. Because only tadpoles of *Lithobates palmipes* and *Corythomantis greeningi* were collected, we used photographs from other localities: Parnamirim municipality, Rio Grande do Norte state, Brazil and Barbalha municipality, Ceará, Brazil, respectively. Adult specimens were preserved in 10% formalin and later stored in 70% alcohol, while tadpoles were preserved and stored in 10% formalin. We used the available literature to confirm the identity of tadpoles collected, when necessary. All specimens collected were deposited at Coleção do Laboratório

de Anfíbios e Répteis (CLAR), Universidade Federal do Rio Grande do Norte (collecting permit 19828-4-ICMBio). Calls were deposited at the “Arquivos Sonoros da Universidade Federal do Rio Grande do Norte” (ASUFRN, the sound archives of Rio Grande do Norte University) collection. Voucher numbers and specimens for loans are available at request.

## Results and Discussion


We recorded 34 species from 15 genera belonging to seven families: Brachycephalidae (1), Bufonidae (2), Cycloramphidae (1), Hylidae (14), Leptodactylidae (13), Microhylidae (2), and Ranidae (1) (Table 1). Several species are reported for the first time in Rio Grande do Norte State (we considered as new record if the species occurrences for the State is not formally published): *Dendropsophus nanus*, *D. soaresi*, *Hypsiboas albomarginatus*, *Ischnocnema ramagii*, *Leptodactylus caatingae*, *L. mystaceus*, *Physalaemus albifrons*, *Scinax cretatus*, *S. fuscomarginatus*, *S. cf. nebulosus* and *S. pachycrus*.




**Figure 2.** Anuran amphibians from Escola Agrícola de Jundiaí, Macaéba municipality, Rio Grande do Norte State, Brazil. A) *Ischnocnema ramagii*, B) *Rhinella granulosa*, C) *R. jimi*, D) *Proceratophrys cristiceps*, E) *Corythomantis greeningi*, F) *Dendropsophus branneri*, G) *D. minutus*, H) *D. oliveirai*.



**Figure 3.** Anuran amphibians from Escola Agrícola de Jundiaí, Macaíba municipality, Rio Grande do Norte State, Brazil. A) *Dendropsophus soaresi*, B) *Hypsiboas albomarginatus*, C) *H. raniceps*, D) *Phyllomedusa nordestina*, E) *Scinax cretatus*, F) *S. fuscomarginatus*, G) *S. cf. nebulosus*, H) *S. pachycrus*.



**Figure 4.** Anuran amphibians from Escola Agrícola de Jundiaí, Macaíba municipality, Rio Grande do Norte State, Brazil. A) *Scinax x-signatus*, B) *Leptodactylus caatingae*, C) *L. fuscus*, D) *L. macrosternum*, E) *L. mystaceus*, F) *L. natalensis*, G) *L. troglodytes*, H) *L. vastus*.



**Figure 5.** Anuran amphibians from Escola Agrícola de Jundiaí, Macaíba municipality, Rio Grande do Norte State, Brazil. A) *Physalaemus albifrons*, B) *P. cuvieri*, C) *Pleurodema diplopeltis*, D) *Pseudopaludicola* cf. *falcipes*, E) *Pseudopaludicola* sp., F) *Elachistoleis cesarii*, G) *Dermatonotus muelleri*, H) *Lithobates palmipes*.

**Table 2.** Number of anuran species found in areas of Atlantic forest and Caatinga domains in north-eastern Brazil (abbreviations for Brazilian states as follows: AL: Alagoas; BA: Bahia; CE: Ceará; PE: Pernambuco; PI: Piauí; SE: Sergipe).

Locality	Species	Duration	Biome	Reference
RPPN Frei Caneca, PE	42	11 months	Atlantic Forest	(Santos and Santos, 2010)
Planalto da Ibiapaba, CE	38	24 months	Caatinga/Moist Forest	(Loebmann and Haddad, 2010)
Macaíba, RN	34	13 months	Atlantic Forest/Caatinga	<b>Present work</b>
Mata do Juncô, SE	33	5 days	Atlantic Forest	(Morato et al., 2011)
Refúgio Ecológico Charles Darwin, PE	30	9 months	Atlantic Forest	(Santos, 2010)
Serra da Jibóia, BA	29	5 months	Atlantic Forest/ Caatinga	(Juncá, 2006)
Aldeia, PE	28	6 months	Atlantic Forest	(Silva and Moura, 2010)
Campo Alegre, AL	28	13 months	Atlantic Forest	(Queissada, 2009)
Reserva Sapiranga, BA	26	5 months	Atlantic Forest/Restinga	(Juncá, 2006)
Parque Nacional Serra de Itabaiana, SE	23	-	Atlantic Forest/Caatinga	(Carvalho et al., 2005)
APA Delta do Parnaíba. PI	21	18 months	Coastal Forest/Caatinga	(Loebmann and Mai, 2008)
Fazenda Maturi, CE	20	1 month	Coastal Forest/Caatinga	(Borges-Nojosa et al., 2010)
Fazenda Formosa, CE	20	1 month	Coastal Forest/Caatinga	(Borges-Nojosa et al., 2010)
Betânia e Floresta, PE	19	7 months	Caatinga	(Borges-Nojosa and Santos, 2005)
Região do Rio Proxim, SE	18	4 months	Atlantic Forest	(Oliveira and Lírio-Junior, 2000)
Floresta do Crasto, SE	17	12 months	Atlantic Forest	(Arzabe et al., 1998)
São José do Bonfim, PB	16	11 months	Caatinga	(Arzabe, 1999)
São João do Cariri, PB	16	23 months	Caatinga	(Vieira et al., 2007)
Mata do Buraqueirinho, PB	14	9 months	Atlantic Forest	(Santana et al., 2008)
Maturéia, PB	12	11 months	Caatinga	(Arzabe, 1999)
Boa Vista, PB	9	23 months	Caatinga	(Vieira et al., 2007)

Among these, *D. nanus*, *D. soaresi*, *L. mystaceus*, *P. albifrons*, *S. fuscomarginatus* and *S. cf. nebulosus* have been reported in the neighbouring Paraíba (Santana, 2010; Santana et al., 2008; Vieira et al., 2007) and Ceará States (Borges-Nojosa et al., 2010; Loebmann and Haddad, 2010). These new records extend species geographic distributions about 150 km to the north of João Pessoa, Paraíba, and 400 km southeast of Pacajús, Ceará. Furthermore, *H. albomarginatus*, *I. ramagii*, *L. caatingae* and *S. cretatus* have their northernmost record in Paraíba State (Santana, 2010; Santana et al., 2008; Vieira et al., 2007), and therefore their geographic distribution is extended approximately 150km to the north.

We only found tadpoles of *Corythomantis greeningi* and *Lithobates palmipes*, in a temporary stream with rocky bottom and a sandy bottom permanent lake, respectively, during the peak of the rainy season. This record extends the known distribution of *C. greeningi* in RN about 150 km east of Angicos, Fazenda São Miguel (Jared et al., 1999), whereas *L. palmipes* has already been reported for regions near Macaíba (Hillis and de-Sá, 1988). This reinforces the importance of tadpole sampling for anuran inventories (Silva, 2010).

The two species of *Pseudopaludicola* are tentatively identified as *P. cf. falcipes*, based on advertisement call, and *Pseudopaludicola* sp. This

genus is considered taxonomically problematic and involves several cryptic species, which are somehow difficult to diagnose (Carvalho, 2012). Likewise, the *Leptodactylus marmoratus* species group complex also has a complicated taxonomy (Angulo et al., 2003) and therefore further analyses are needed to confirm the identification of this species.

We collected species typical of the Atlantic Forest morphoclimatic domain, such as *D. branneri*, *H. albomarginatus*, *I. ramagii* and *S. cretatus* (Nunes and Pombal, 2011; Santana et al., 2008) and Caatinga domain, such as *C. greeningi*, *P. albifrons*, *Pleurodema diplolister*, and *Phyllomedusa nordestina* (Arzabe, 1999; Caramaschi, 2006; Rodrigues, 2003). This reinforces the transitional nature of our study site. Accordingly, Cestaro and Soares (2004) found a similar result for plants and characterized the area as a transition region between the Caatinga and Atlantic Forest phytogeographic provinces.

Because of the lack of publications on amphibian communities from RN, it is not possible to compare EAJ species richness with other localities in the state. However, even with different methodologies (specially field effort, area size, and sampling methods), we found a high species richness in comparison to other studies conducted in the Atlantic Forest and Caatinga in north-eastern Brazil (Table 2). This supports the hypothesis

that transition areas may be especially important for biodiversity management and conservation (Remanamanjato et al., 2002). Hence, prioritization of such areas of ecological tension can help protect biodiversity of neighbouring biomes, especially in regions where human activity has devastated much of the original vegetation, as is the case of RN.

**Acknowledgements.** We thank Bruno França and Tonny Marques for help during fieldwork. We also thank Ulisses Caramaschi, Ronald Heyer, Gindomar Gomes Santana, and Paulo Garcia for help with adult and tadpole identifications. Bruno França and Felipe Figueirêdo kindly shared some of their pictures presented in this paper. Parts of this research were financed by a grant from FAPERN (“Programa Primeiros Projetos–PPP”) awarded to AAG.

## References

- Ab’Saber, A.N. (1977): Os domínios morfoclimáticos na América do Sul. *Geomorfologia* (São Paulo) **52**: 1–23.
- Angulo, A., Crocroft, R.B., Reichle, S. (2003): Species identity in the genus *Adenomera* (Anura:Leptodactylidae) in southeastern Peru. *Herpetologica* **59**(4): 490–504.
- Arzabe, C. (1999): Reproductive activity patterns of anurans in two different altitudinal sites within the Brazilian Caatinga. *Rev. Bras. Zool.* **16**(3): 851–864.
- Arzabe, C., Carvalho, C.X., Costa, M.A.G. (1998): Anuran assemblages in Crasto forest ponds (Sergipe State, Brazil): Comparative structure and calling activity patterns. *Herpetol. J.* **8**: 111–113.
- Borges-Nojosa, D.M., Cascon, P. (2005): Herpetofauna da Área Reserva da Serra das Almas, Ceará. In: *Análise das Variações da Biodiversidade do Bioma Caatinga*, p. 245–260. Araújo, F.S., Rodal, M.J.N., Barbosa, M.R.V., Eds., Brasília, Ministério do Meio Ambiente.
- Borges-Nojosa, D.M., Prado, F.M.V., Borges-Leite, M.J., Gurgel-Filho, N.M., Bacalini, P. (2010): Avaliação do Impacto do Manejo Florestal Sustentável na Herpetofauna de Duas Áreas de Caatinga nos Municípios de Caucaia e Pacajus no Estado do Ceará. In: *Uso Sustentável e Conservação dos Recursos Florestais da Caatinga*, p. 315–330. Gariglio, M.A., Sampaio, E.V.S.B., Cestaro, L.A., Kageyama, P.Y., Eds., Brasília, Serviço Florestal Brasileiro: MMA.
- Borges-Nojosa, D.M., Santos, E.M. (2005): Herpetofauna da Área de Betânia e Floresta, Pernambuco. In: *Análise das Variações da Biodiversidade do Bioma Caatinga*, p. 275–289. Araújo, F.S., Rodal, M.J.N., Barbosa, M.R.V., Eds., Brasília, Ministério do Meio Ambiente.
- Camardelli, M., Napoli, M.F. (2012): Amphibian conservation in the Caatinga Biome and Semiarid region. *Herpetologica* **68**(1): 31–47.
- Caramaschi, U. (2006): Redefinição do grupo de *Phyllomedusa hypochondrialis*, com redescrição de *P. megacephala* (Miranda-Ribeiro, 1926), revalidação de *P. azurea* Cope, 1862 e descrição de uma nova espécie (Amphibia, Anura, Hylidae). *Arq. Mus. Naci.* **64**(2): 159–179.
- Carvalho, C.M., Villar, J.C., Oliveira, F.F. (2005): Répteis e anfíbios. In: *Parque Nacional Serra de Itabaiana: Levantamento da Biota*, p. 39–61. Carvalho, C.M., Villar, J.C., Eds., Aracaju, IBAMA.
- Carvalho, T.R. (2012): A new species of *Pseudopaludicola* Miranda-Ribeiro (Leiuperinae: Leptodactylidae: Anura) from the Cerrado of southeastern Brazil with a distinctive advertisement call pattern. *Zootaxa* **3328**: 47–54.
- Cestaro, L.A., Soares, J.J. (2004): Variações florística e estrutural e relações fitogeográficas de um fragmento de floresta decídua no Rio Grande do Norte, Brasil. *Acta Bot. Bras.* **18**(2): 203–218.
- Cushman, S.A. (2006): Effects of habitat loss and fragmentation on amphibians: A review and prospectus. *Biodivers. Conserv.* **128**: 231–240.
- Duellman, W.E. (1999): *Patterns of Distribution of Amphibians: A Global Perspective*. Baltimore, The Johns Hopkins University Press.
- Garda, A.A., São-Pedro, V.A., Lion, M.B. (2010): The advertisement and release calls of *Rhinella jimi* (Anura, Bufonidae). *S. Am. J. Herpetol.* **5**(2): 151–156.
- Hillis, D.M., de-Sá, R.O. (1988): Phylogeny and taxonomy of the *Rana palmipes* group (Salientia: Ranidae). *Herpetol. Monogr.* **2**: 1–26.
- Jared, C., Antoniazzi, M.M., Katchburian, E., Toledo, R.C., Freymuller, E. (1999): Some aspects of the natural history of the casque-headed tree frog *Corythomantis greeningi* Boulenger (Hylidae). *Ann. Sci. Nat.* **3**: 105–115.
- Juncá, F.A. (2006): Diversidade e uso de hábitat por anfíbios anuros em duas localidades de Mata Atlântica, no norte do estado da Bahia. *Biota Neotrop.* **6**(2): 1–17.
- Loebmann, D., Haddad, C.F.B. (2010): Amphibians and reptiles from a highly diverse area of the Caatinga domain: composition and conservation implications. *Biota Neotrop.* **10**(3): 227–256.
- Loebmann, D., Mai, A.C.G. (2008): Amphibia, Anura, Coastal Zone, State of Piauí, Northeastern Brazil. Check List **4**(2): 161–170.
- Magalhães, F.M., Santana, D.J., Neto, A.M., Garda, A.A. (2012): The tadpole of *Elachistocleis cesarii* Miranda-Ribeiro, 1920 (Anura, Microhylidae). *Zootaxa* **3187**: 54–56.
- Mittermeier, R., A., Myers, N., Gil, P.R., Mittermeier, C.G. (1999): Hotspots: Earth’s Biologically Richest and Most Endangered Terrestrial Ecoregions. Mexico City, CEMEX Conservation International.
- Morato, S.A.A., Lima, A.M.X., Staut, D.C.P., Faria, R.G., Souza-Alves, J.P., Gouveia, S.F., Scupino, M.R.C., Gomes, R., Silva, M.J. (2011): Amphibians and Reptiles of the Refúgio de Vida Silvestre Mata do Junco, municipality of Capela, state of Sergipe, northeastern Brazil. Check List **7**(6): 756–762.
- Moura, G.J.B. (2010): Estrutura da Comunidade de Anuros e Lagartos de Remanescente de Mata Atlântica, Com Considerações Ecológicas e Zoogeográficas Sobre a Herpetofauna do Estado de Pernambuco. PhD Dissertation, Universidade Federal da Paraíba, João Pessoa.

- Nunes, I., Pombal Jr., J.P. (2011): A new snouted treefrog of the speciose genus *Scinax* Wagler (Anura, Hylidae) from north-eastern Brazil. *Herpetologica* **67**(1): 80–88.
- Oliveira, F.F., Lírio-Junior, G.P. (2000): Anfíbios anuros do campus da Universidade Federal de Sergipe. *Biol. Ger. Exper.* **1**(1): 42–74.
- Queissada, I.C.S.T. (2009): Diversidade da herpetofauna de uma área de Mata Atlântica do Estado de Alagoas: A reserva particular da usina Porto Rico, Campo Alegre. PhD Dissertation, Universidade Estadual Paulista, Rio Claro.
- Remanamanjato, J., McIntyre, P.B., Nussbaum, R.A. (2002): Reptile, amphibian, and lemur diversity of the Malahelo Forest, a biogeographical transition zone in southeastern Madagascar. *Biodivers. Conserv.* **11**: 1791–1807.
- Rodrigues, M.T. (2003): Herpetofauna da Caatinga. In: Biodiversidade, ecologia e conservação da Caatinga, p. 181–236. Tabarelli, M., Silva, J.M.C., Eds., Recife, Brasil., Universidade Federal de Pernambuco.
- Santana, D.J., Mesquita, D.O., Garda, A.A. (2011): The advertisement call of *Dendropsophus oliveirai* (Anura, Hylidae). *Zootaxa* **2997**: 67–68.
- Santana, G.G. (2010): Taxocenose de anfíbios anuros habitando um remanescente de Floresta Atlântica (Zona da Mata do estado da Paraíba, Nordeste do Brasil): composição de espécies, utilização dos habitats e disponibilidade de recursos alimentares. PhD Dissertation, Universidade Federal da Paraíba, João Pessoa.
- Santana, G.G., Vieira, W.L.S., Lima, Y.C.C., Vieira, K.S. (2008): Herpetofauna em um fragmento de Floresta Atlântica no estado da Paraíba, Região Nordeste do Brasil. *Biotaemas* **21**: 75–84.
- Santos, E.M. (2010): Anfíbios anuros do Refúgio Ecológico Charles Darwin, Igarassu, Pernambuco, Brasil. In: Herpetofauna no Estado de Pernambuco, p. 124–132. Moura, G.J.B., Santos, E.M., Oliveira, M.A.B., Cabral, M.C.C., Eds., Brasília, Ministério do Meio Ambiente.
- Santos, S.P.L., Santos, E.M. (2010): Anurofauna da Reserva Particular do Patrimônio Natural Frei Caneca, Município de Jaqueira, Estado de Pernambuco, Brasil. In: Herpetofauna no Estado de Pernambuco, p. 184–196. Moura, G.J.B., Santos, E.M., Oliveira, M.A.B., Cabral, M.C.C., Eds., Brasília, Ministério do Meio Ambiente.
- São-Pedro, V.A., Medeiros, P.H., Garda, A.A. (2011): The advertisement call of *Rhinella granulosa* (Anura, Bufonidae). *Zootaxa* **3092**: 60–62.
- Segalla, M.V., Caramaschi, U., Cruz, C.A.G., Garcia, P.C.A., Grant, T., Haddad, C.F.B., Langone, J. (2012): Brazilian amphibians – List of species. Accessible at <http://www.sbhherpetologia.org.br>. Sociedade Brasileira de Herpetologia (Accessed on June 2012).
- Silva, F.R. (2010): Evaluation of survey methods for sampling anuran species richness in the neotropics. *S. Am. J. Herpetol.* **5**(3): 212–220.
- Silva, R.G., Moura, G.J.B. (2010): Abundância, riqueza e especificidade microambiental da anurofauna no complexo da Mata Atlântica de Aldeia (PE). In: Herpetofauna no Estado de Pernambuco, p. 134–146. Moura, G.J.B., Santos, E.M., Oliveira, M.A.B., Cabral, M.C.C., Eds., Brasília, Ministério do Meio Ambiente.
- Tabarelli, M., Silva, J.M.C. (2003): Áreas e ação prioritárias para a conservação da Caatinga. In: Biodiversidade, ecologia e conservação da Caatinga, p. 781–800. Leal, I., Silva, J.M.C., Tabarelli, M., Eds., Recife, Brasil, Universidade Federal de Pernambuco.
- Vieira, W.L.S., Arzabe, C., Santana, G.G. (2007): Composição e distribuição espaço-temporal de anuros no Cariri paraibano, Nordeste do Brasil. *Oecol. Brasil.* **11**(3): 383–396.
- Willians, P.H. (1996): Mapping variations in strength and breadth of biogeographic transition zones using species turnover. *P. R. Soc. London* **263**(1370): 579–588.