Predation attempt on *Trachycephalus typhonius* (Linnaeus, 1758) by *Caiman yacare* (Daudin, 1802)

Kelvin Y. Mori^{1,*}, Sean Keuroghlian-Eaton^{1,2}, Diego G. Cavalheri^{1,2}, João E. de Almeida-Júnior¹, Márcia M. P. Müller^{1,2}, and Diego J. Santana¹

Frogs are an important resource for many animal groups, such as fish, mammals, birds, reptiles, and invertebrates (Toledo et al., 2007; Wells, 2007). To avoid predation, frogs have developed a variety of defensive behaviours and mechanisms, including puffing up their bodies, biting, and producing toxic secretions (Ferreira et al., 2019). The genus *Trachycephalus* is widely distributed in the neotropical region (Frost, 2021) and secretes a large amount of toxins with adhesive effects on their dorsum (e.g., Toledo et al., 2011; Ferreira et al., 2019).

Trachycephalus typhonius (Linnaeus, 1758) occurs from southern Mexico to northern Argentina (Leenders, 2016; Piva, 2020), and is a common species in the Pantanal Floodplain. Although this large frog is mainly arboreal, it reproduces in shallow temporary or permanent ponds during the wet season (Leenders, 2016). The Yacare Caiman (*Caiman yacare*), another common species in Pantanal floodplains, is an opportunistic predator that feeds on a variety of animals depending on availability (Santos et al., 1996). In this study, we report an attempted predation by a juvenile *C. yacare* on an adult *T. typhonius* in the Pantanal Floodplain, Mato Grosso do Sul, Brazil.

On 14th December 2021 at around 19:29 h, we observed a juvenile *C. yacare* with an adult *T. typhonius* in its mouth at the edge of a temporary pond (Fig. 1), during fieldwork at the Base de Estudos do Pantanal in the municipality of Corumbá, state of Mato Grosso do Sul, Brazil (-19.577°N, -57.019°E). The event was

recorded using a video camera (Panasonic HC-V270), and the video has been deposited in the Zoological Collection of the Universidade Federal de Mato Grosso do Sul - Digital Media (ZUFMS-MID) under the vouchers ZUFMS-MID00005 and ZUFMS-MID00006. We observed the event for 10 minutes until the C. yacare released its prey and dove underwater swimming away, not appearing again. At the time of the event, 10 species of anurans were actively calling in the swamp, including Trachycephalus typhonius, Rhinella diptycha, Elachistocleis bicolor, Physalaemus albonotatus, Scinax acuminatus, Dendropsophus nanus, Rhinella bergi, Leptodactylus podicipinus, Leptodactylus fuscus, and Leptodactvlus macrosternum. We counted at least eight caimans, all juveniles, which around 21:00 h began to leave the swamp and move to the surrounding bushes.

Caiman yacare is known to occasionally include anurans in its diet, but they are not considered a significant part of it (Santos et al., 1996; Toledo et al., 2007). This could be due to the fast digestion rate of amphibians, which makes them less likely to be detected using stomach flushing as a diet assessment method (Delany and Abercrombie, 1986; Platt et al., 2006). The diet of caimans depends on prey availability (Magnusson et al., 1987; Santos et al., 1996; da Silveira and Magnusson, 1999), and during the rainy season in the Pantanal, the anuran population significantly increases (Alho, 2008). In a study on Caiman crocodilus, the sister species of C. yacare, in Venezuela during the wet season, Gorzula (1978) found that 66% of the prey taken were anurans. Bernal (2006) reported that adult C. crocodilus responded to playback of the advertisement calls of Leptodactylus pentadactylus and Rhinella marina, indicating that predation on anurans by crocodilians may be more common and that frogs could represent a more significant part of caiman's diet than previously thought. Therefore, further dietary studies on C. yacare are needed to better understand the importance of anurans in the diet of this species.

¹ Mapinguari - Laboratório de Biogeografia e Sistemática de Anfibios e Répteis, Universidade Federal de Mato Grosso do Sul, Avenida Costa e Silva S/N - Pioneiros, Campo Grande, 79002-970 Mato Grosso do Sul, Brazil.

² Universidade Estadual Paulista (UNESP), São José Do Rio Preto, São Paulo 15054-000, Brazil.

^{*} Corresponding author. E-mail: kelvinyuiti@gmail.com

^{© 2023} by Herpetology Notes. Open Access by CC BY-NC-ND 4.0.



Figure 1. Attempted predation by a juvenile *Caiman yacare* on an adult *Trachycephalus typhonius* at the edge of a temporary pond in Base de Estudos do Pantanal, Corumbá Municipality, Mato Grosso do Sul, Brazil. Photographs by Kelvin Yuiti Mori (A) and Sean Keuroghlian-Eaton (B).

Acknowledgments. We are thankful to PELD Pantanal (Programa de Pesquisa Ecológica de Longa Duração), Fundect (Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e Tecnologia de Mato Grosso do Sul), Semagro (Secretaria de Meio Ambiente, Desenvolvimento Econômico, Produção e Agricultura Familiar), UFMS (Universidade Federal do Mato Grosso do Sul), ZUFMS (Coleção Zoológica de Referência da Universidade Federal de Mato Grosso do Sul) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for all the support. DJS thanks Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq proceses 404239/2021-8, 402519/2016-7) for financial support. DJS also thanks CNPq for his research fellowship (CNPq 309420/2020-2).

References

- Alho, C.J.R. (2008): Biodiversity of the Pantanal: Response to seasonal flooding regime and to environmental degradation. Brazilian Journal of Biology 68: 957–966.
- Bernal, X.E. (2006): Caiman crocodilus (spectacled caiman). Predation/eavesdropper. Herpetological Review 37: 460–461.

- Carrillo, J.F.C. (2019): *Caiman yacare* (yacare caiman, jacaré do pantanal). Tadpoles in diet. Herpetological Review 50: 130– 131.
- Da Silveira, R., Magnusson, W.E. (1999): Diets of spectacled and black Caiman in the Anavilhanas Archipelago, Central Amazonia, Brazil. Journal of Herpetology 33(2): 181–192.
- Delany, M.F., Abercrombie, C.L. (1986): American Alligator Food Habits in Northcentral Florida. The Journal of Wildlife Management 50(2): 348.
- Ferreira, R.B., Lourenço-de-Moraes, R., Zocca, C., Duca, C., Beard, K.H., Brodie, E.D. (2019): Antipredator mechanisms of post-metamorphic anurans: a global database and classification system. Behavioral Ecology and Sociobiology 73: 69.
- Frost, D.R. (2021): Amphibian Species of the World: An Online Reference. Version 6.1. American Museum of Natural History, New York, New York, USA. Available at: https:// amphibiansoftheworld.amnh.org/Amphibia/Anura/Hylidae/ Lophyohylinae/Trachycephalus. Accessed on 2 June 2023.
- Gorzula, S.J. (1978): An ecological study of *Caiman crocodilus crocodilus* inhabiting savanna lagoons in the Venezuelan Guayana. Oecologia 35(1): 21–34.

- Leenders, T. (2016): Amphibians of Costa Rica: a field guide. Ithaca, USA, London, ENG, Cornell University Press. 446p.
- Magnusson, W.E., da Silva, E.V., Lima, A.P. (1987): Diets of Amazonian crocodilians. Journal of Herpetology 21(2): 85–95.
- Piva, A., dos Santos, F.M., da Silva Costa-Urquiza, A., Soares, M.P., Urquiza, M.V.S., de Albuquerque, N.R. (2020): Anurans of the parque municipal de piraputangas, on the western border of the Pantanal, Mato Grosso do Sul, Brazil. Check List 16(6): 1709–1724.
- Platt, S.G., Rainwater, T.R., Finger, A.G., Thorbjarnarson, J.B., Anderson, T.A., McMurry, S.T. (2006): Food habits, ontogenetic dietary partitioning and observations of foraging behaviour of Morelet's crocodile (*Crocodylus moreletii*) in northern Belize. Herpetological Journal 16(3): 281–290.
- Santos, S.A., Nogueira, M.S., Pinheiro, M.S., Campos, Z., Magnusson, W.E., Mourao, G.M. (1996): Diets of *Caiman crocodilus yacare* from different habitats in the Brazilian Pantanal. Herpetological Journal 6(4): 111–117.
- Toledo, L.F., Ribeiro, R.S., Haddad, C.F.B. (2007): Anurans as prey: An exploratory analysis and size relationships between predators and their prey. Journal of Zoology 271(2): 170–177.
- Toledo, L.F., Sazima, I., Haddad, C.F.B. (2011): Behavioural defences of anurans: An overview. Ethology Ecology and Evolution 23(1): 1–25.
- Wells, K.D. (2007): The Ecology and Behavior of Amphibians. In The Ecology and Behavior of Amphibians.