

Distribution extension, geographic distribution maps and comments on the advertisement call of *Pseudopaludicola pocoto* Magalhães, Loebmann, Kokubum, Haddad and Garda, 2014 (Anura: Leptodactylidae: Leiuperinae)

Elvis Almeida Pereira^{1,2,*}, Henrique Folly^{2,3}, Raoni Rebouças¹, and Diego José Santana²

The leptodactylid genus *Pseudopaludicola* comprises 21 species of small frogs occurring from northern Argentina to Venezuela, associated with open areas across Neotropical forests and open formations (e.g., Chaco, Pampas, Cerrado, and Caatinga) east of the Andes (Carvalho, 2012; Pansonato et al., 2012; Andrade et al., 2016; Frost, 2017). Several *Pseudopaludicola* species have been described recently, increasing considerably our knowledge about this genus (Toledo, 2010; Toledo et al., 2010; Carvalho, 2012; Pansonato et al., 2012; Roberto et al., 2013; Magalhães et al., 2014). Nevertheless, recognition of these species based on external morphology only has become harder (Roberto et al., 2013), as a result, most species of this genus are similar and diagnosed based mainly through differences on their advertisement calls (Carvalho, 2012; Roberto et al., 2013; Magalhães et al., 2014).

Pseudopaludicola pocoto Magalhães, Loebmann, Kokubum, Haddad and Garda, 2014 was described from the Municipality of Santa Quitéria, State of Ceará, Brazil. It was also recorded in the states of Rio Grande do Norte, Paraíba and Pernambuco, in northeastern

Brazil, in areas associated with the Caatinga Biome (Magalhães et al., 2014; Santana et al., 2015; Lantyer-Silva et al., 2016). Recently, Andrade et al. (2015, 2017) extended the species' distribution to the states of Bahia and Minas Gerais. The record by Andrade et al. (2015) was the first for the species in the Atlantic Forest, in southeastern Brazil. However, this new record, caused a large distribution gap, because about 800 km separates the records between the municipalities of Xique-Xique in the state of Bahia (Andrade et al., 2017) and Carlos Chagas in the state of Minas Gerais (Andrade et al., 2015).

Herein, we present two new records for this species; a second record of *P. pocoto* in the state of Bahia, and the second record in the Atlantic Forest Biome, of the



Figure 1. Individual of *Pseudopaludicola pocoto* registered in the Municipality of Catuji, Minas Gerais state.

¹ Universidade Federal Rural do Rio de Janeiro (UFRRJ), Departamento de Biologia Animal, Laboratório de Herpetologia, CEP 23890-000 Seropédica, Rio de Janeiro, Brazil.

² Matinguari - Laboratório de Biogeografia e Sistemática de Anfíbios e Répteis, Universidade Federal de Mato Grosso do Sul, 79002-970 Campo Grande, Mato Grosso, Brazil.

³ Programa de Pós-graduação em Biologia Animal, Departamento de Biologia Animal, Universidade Federal de Viçosa, 36570-000 Viçosa, Minas Gerais, Brazil.

* Corresponding author. E-mail: elvisaps@hotmail.com




Figure 2. Map showing the known geographic distribution of *Pseudopaludicola pocoto*. Literature records (circles) and new records (triangles).

state of Minas Gerais (Fig. 1), and provide comments on its advertisement call. We collected one adult male of *P. pocoto* on 12 November 2014, in the Municipality of Caetité, State of Bahia, Brazil (-14.070363°S, -42.484881°W, WGS84, 907 m a.s.l.) and four adult males on 29 May 2016, in the Municipality of Catuji, State of Minas Gerais (-17.391400°S, -41.490371°W, WGS84, 395 m a.s.l.). We recorded the advertisement call of one individual from Municipality Caetité and one from Municipality Catuji and, compared it to the calls previously described for the species (Magalhães *et al.*, 2014; Andrade *et al.*, 2015, 2017). The advertisement calls was recorded using a Sony IC Recorder ICD-PX312F with an internal microphone, MP3 File, 48 kHz, 16 bits, while the individuals were calling in a small puddle of water on sand. We analysed the calls with Raven Pro 1.5, 64-bit version (Bioacoustics Research Program 2014) with the following settings: window type = Hann, window size = 256 samples, 3 dB filter bandwidth = 248 or 270 Hz, brightness = 50%, contrast = 50%, overlap = 75% (locked), hop size = 38 samples,

DFT size = 1024 samples (locked), grid spacing = 43.1 or 46.9 Hz, frame = 100, flat top filter and generated audio spectrograms in R software using the package “Seewave” (Sueur *et al.*, 2008; R Core Team, 2016). We followed the same characteristics and terminologies for calls descriptions adopted by Magalhães *et al.* (2014). All collected specimens are housed in the Coleção Zoológica de Referência da Universidade Federal de Mato Grosso do Sul (ZUFMS), under the following numbers: ZUFMS-AMP07451 (snout-vent length - SVL = 12.72 mm) and ZUFMS-AMP07452 (SVL = 13.99 mm).

The new record in Caetité extend the known distribution in approximately 360 km southward from Xique-Xique, State of Bahia (Andrade *et al.*, 2017) and 450 km northward from Carlos Chagas, State of Minas Gerais (Andrade *et al.*, 2015), contributing to extension of distribution within the Caatinga Biome. However, the record from Catuji, State of Minas Gerais, constitutes the second for the Atlantic Forest Biome. These records provided a significant increase in the distribution for




Figure 3. Spectrogram (top), and amplitude of the waveform (bottom) from a two call sequence of *Pseudopaludicola pocoto* advertisement call recorded at Municipality of Caetité, State of Bahia, Brazil (SVL = 13.99 mm).

this species and bridged the gap on our knowledge of its distribution (Fig. 2).

Many species of frogs are typical of the Caatinga Biome and the northern portion of the Atlantic Forest Biome, such as *Boana albomarginata* (Spix, 1824) (Magalhães et al., 2013), *Corythomantis greeningi* Boulenger, 1896 (Godinho et al., 2013; Silva et al., 2014); *Leptodactylus caatingae* Heyer and Juncá, 2003 (Heyer and Juncá, 2003; Vieira et al., 2006); *Physalaemus cicada* Bokermann, 1966 (Lisboa and Haddad, 2009);

Pipa carvalhoi (Miranda-Ribeiro, 1937) (Santana et al., 2014), *Pleurodema diplolister* (Peters, 1870) (Andrade and Vaz-Silva, 2012); *Proceratophrys cristiceps* and *Rhinella granulosa* (Cassemiro et al., 2012). Thus, these two new records in the Atlantic Forest biome may indicate that *P. pocoto* also has a broad distribution along these biomes.

Pseudopaludicola pocoto was previously known to occur only in sites of the Caatinga Biome or ecotone sites (Magalhães et al., 2014), but here the two records




Figure 4. Spectrogram (top), and amplitude of the waveform (bottom) from a two call sequence of *Pseudopaludicola pocoto* advertisement call recorded at Municipality of Catuji, State of Minas Gerais, Brazil (SVL = 12.72 mm).

Table 1. Advertisement call traits of *Pseudopaludicola pocoto* from Municipality of Caetité, Bahia state (BA), Brazil (present study), and Catuji, Minas Gerais state (MG), Brazil (present study) and comparative data for populations from Northeastern Brazil (Magalhães *et al.*, 2014), Municipality of Carlos Chagas, Minas Gerais state, Brazil (Andrade *et al.*, 2015), Municipality of Cabrobó, Pernambuco state, Brazil (Lantyer-Silva *et al.*, 2016) and Municipality of Xique-Xique, Bahia state, Brazil (Andrade *et al.*, 2017). Mean \pm SD (minimum–maximum); n = number of males recorded (number of analysed notes).

Acoustic variables	Caetité-BA (present study) $n = 1$ (40)	Catuji-MG (present study) $n = 1$ (14)	Magalhães <i>et al.</i> (2014) $n = 9$ (135)	Andrade <i>et al.</i> (2015) $n = 2$ (30)	Lantyer-Silva <i>et al.</i> (2016) $n = 1$ (30)	Andrade <i>et al.</i> (2017) $n = 5$ (50)
Note duration (ms)	263 \pm 15 (233–291)	306 \pm 19 (264–326)	238 \pm 31 (126–290)	265 \pm 32 (205–323)	274 \pm 22 (223–309)	322 \pm 36 (170–397)
Internote interval (ms)	230 \pm 37 (186–411)	202 \pm 19 (181–250)	220 \pm 59 (136–507)	251 \pm 61 (196–447)	203 \pm 23 (181–263)	251 \pm 15 (209–291)
Notes per minute	–	–	124 \pm 16 (94–150)	–	–	108 \pm 6.5 (100–118)
Pulses per note	3.0 \pm 0.5 (2–3)	3	3	3	–	3.0 \pm 0.05 (2–3)
Pulse duration (ms)	9 \pm 1 (6–14)	8 \pm 1 (6–10)	5 \pm 1 (3–8)	6 \pm 1 (3–8)	12 \pm 2 (7–18)	8 \pm 1 (6–10)
Interpulse interval (ms)	137 \pm 65 (12–241)	137 \pm 11 (105–154)	111 \pm 21 (43–166)	122 \pm 31 (54–205)	138 \pm 40 (82–263)	151 \pm 17 (112–187)
Pulse rate (pulses/second)	10.8 \pm 1.5 (7.6–12.8)	9.7 \pm 0.01 (9.2–11.3)	13 \pm 2 (10–18)	11 \pm 2 (8–15)	–	9.4 \pm 1 (7.6–12)
Dominant frequency (Hz)	5340.2 \pm 0 (5340.2)	5684.8 \pm 85.6 (5684.8–5657)	5636 \pm 300 (5168–6374)	5001 \pm 90 (4823–5168)	5804 \pm 107 (5625–6000)	5133 \pm 269.8 (4734–5437)
Air Temperature (°C)	–	–	24–25	20	–	27

in Minas Gerais are within the Atlantic Forest and not in ecotone areas with Cerrado or Caatinga. Despite these new distributional information, there are still gaps along the species distribution, and further studies are needed to elucidate how wide this species distribution actually is.

The call of *P. pocoto* (Figs. 3, 4) is characterized by a single note emitted as a series of well-defined sequences of two to three pulses and they overlap in almost all acoustic parameters analysed by other authors, except the dominant frequency, which was higher in the individual of Caetité, Bahia (=5340.2 Hz) than that described for Carlos Chagas, Minas Gerais (=5001 \pm 90 Hz) (Andrade *et al.*, 2015) and lower than that of Cabrobó, Pernambuco (=5804 \pm 107 Hz) (Lantyer-Silva *et al.*, 2016). The calls of the individuals from Catuji-Minas Gerais (=5684.8 \pm 85.6 Hz) had the dominant frequency higher than those of Carlos Chagas (=5001 \pm 90 Hz) and Xique-Xique-Bahia (=5133 \pm 269.8 Hz) (Andrade *et al.*, 2015, 2017) (Table 1). The calls analysed here, as in Andrade *et al.* (2017), have an amplitude increase between the first and second pulse, followed by a

decrease in amplitude between the second and third pulse. However, the dominant frequency remained the same between the three pulses, different from those analysed by Magalhães *et al.* (2014) and Andrade *et al.* (2015) in which the calls exhibited upward frequency modulation, and in Andrade *et al.* (2017), the calls had a higher frequency in the second pulse.

Overall, the call of *P. pocoto* is similar along the entire species distribution. As data on their distributions are still deficient (Azevedo-Ramos and Galatti, 2002; Silvano and Segalla, 2005) it is possible that the distribution of *P. pocoto* could be larger and/or highly fragmented (Andrade *et al.*, 2015). Further, more studies are needed in order to increase our knowledge of this species distribution.

Acknowledgments. We are grateful to Hélio Ricardo and Felipe Andrade for suggestions on the manuscript. EAP and HF are grateful to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for their fellowships, respectively. EAP thanks The Cornell Lab of Ornithology (Bioacoustics Research Program) for the license for Raven Pro:

Interactive Sound Analysis Software and Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio/SISBIO for the license collection.

References

- Andrade, S.P., Vaz-Silva, W. (2012): First state record and distribution extension of *Pleurodema dipolister* (Peters, 1870) (Anura: Leiuperidae) from State of Goiás, Brazil. *Check List* **8**: 149–151.
- Andrade, I.S., Barros, L.C.S., Oliveira, A.F.S., Juncá, F.A., Magalhães, F.M. (2015): Distribution extension of *Pseudopaludicola pocoto* Magalhães, Loebmann, Kokubum, Haddad and Garda, 2014 (Anura: Leptodactylidae: Leiuperinae) in State of Minas Gerais, Brazil. *Herpetology Notes* **8**: 227–229. <http://biotaxa.org/hn/article/view/9474>.
- Andrade, F.S., Magalhães, F.M., Nunes-de-Almeida, C.H.L., Veiga-Menoncello, A.C.P., Santana, D.J., Garda, A.A., et al. (2016): A new species of long-legged *Pseudopaludicola* from northeastern Brazil (Anura, Leptodactylidae, Leiuperinae). *Salamandra* **52**: 107–124.
- Andrade, F.S., Leite, F.S.F., Carvalho, T.R., Bernardes, C.S., Giaretta, A.A. (2017): First record of *Pseudopaludicola pocoto* Magalhães, Loebmann, Kokubum, Haddad and Garda, 2014 (Anura, Leptodactylidae, Leiuperinae) in Bahia state, Northeastern Brazil, with further data on its advertisement call. *Check List* **13**: 2047 doi: <https://doi.org/10.15560/13.1.2047>.
- Azevedo-Ramos, C., Galatti, U. (2002): Patterns of amphibian diversity in Brazilian Amazônia: conservation implications. *Biological Conservation* **103**: 103–111.
- Bioacoustics Research Program. (2014): Raven Pro: Interactive Sound Analysis Software, version 1.5. The Cornell Lab of Ornithology, Ithaca, New York. Available at <http://www.birds.cornell.edu/raven>. Accessed on 27 March 2016.
- Carvalho, T.R. (2012): A new species of *Pseudopaludicola* Miranda-Ribeiro (Leiuperinae: Leptodactylidae: Anura) from the Cerrado of southeastern Brazil with a distinctive advertisement call pattern. *Zootaxa* **3328**: 47–54.
- Cassemiro, F.A.S., Gouveia, S.F., Diniz-Filho, J.A.F. (2012): Distribuição de *Rhinella granulosa*: integrando envelopes bioclimáticos e respostas ecofisiológicas. *Revista da Biologia* **8**: 38–44.
- Frost, D.R. (2017): Amphibian species of the world: an online reference. Version 6.0. American Museum of Natural History. Available at <http://research.amnh.org/vz/herpetology/amphibia>. Accessed on 21 March 2017.
- Godinho, L.B., Moura, M.R., Feio, R.N. (2013): New records and geographic distribution of *Corythomantis greeningi* Boulenger, 1896 (Amphibia: Hylidae). *Check List* **9**: 148–150.
- Heyer, W.R., Juncá, F.A. (2003): *Leptodactylus caatingae*, a new species of frog from eastern Brazil (Amphibia: Anura: Leptodactylidae). *Proceedings of the Biological Society of Washington* **116**: 317–239.
- Lantyer-Silva, A.S.F., Matos, M.A., Gogliath, M., Marciano-Jr, E., Nicola, P.A. (2016): New records of *Pseudopaludicola pocoto* Magalhães, Loebmann, Kokubum, Haddad and Garda, 2014 (Amphibia: Anura: Leptodactylidae) in the Caatinga Biome, Brazil. *Check list* **12**: 1–4.
- Lisboa, B.S., Haddad, C.F.B. (2009): Amphibia, Anura, Leiuperidae, *Physalaemus cicada* Bokermann, 1966: Distribution extension and geographic distribution map. *Check List* **5**: 699–701.
- Magalhães, F.M., Dantas, A.K.B.P., Brito, M.R.M., Medeiros, P.H.S., Oliveira, A.F., Pereira, T.C.S.O., et al. (2013): Anurans from an Atlantic Forest-Caatinga ecotone in Rio Grande do Norte State, Brazil. *Herpetology Notes* **6**: 1–10.
- Magalhães, F.M., Loebmann, D., Kokubum, M.N.C., Haddad, C.F.B., Garda, A.A. (2014): A new species of *Pseudopaludicola* (Anura: Leptodactylidae: Leiuperinae) from northeastern Brazil. *Herpetologica* **70**: 77–88.
- Pansonato, A., Morais, D.H., Avila, R.W., Kawashita-Ribeiro, R.A., Strussmann, C., Martins, I.A. (2012): A new species of *Pseudopaludicola* Miranda-Ribeiro, 1926 (Anura: Leiuperidae) from the State of Mato Grosso, Brazil, with comments on the geographic distribution of *Pseudopaludicola canga* Giaretta and Kokubum, 2003. *Zootaxa* **3523**: 49–58.
- R Core Team (2016): R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Version 3.1.2. Available at <http://www.R-project.org>. Accessed on 29 March 2017.
- Roberto, I.J., Cardozo, D. Avila, R.W. (2013): A new species of *Pseudopaludicola* (Anura, Leiuperidae) from western Piauí state, Northeast Brazil. *Zootaxa* **3636**: 348–360.
- Santana, D. O., Franco, S. da C., Rocha, S. M., Freitas, E. B. de, De-Carvalho, C. B., Faria, R. G. (2014): First record of *Pipa carvalhoi* (Miranda-Ribeiro, 1937) (Anura: Pipidae) in the State of Sergipe, Northeastern Brazil. *Check List. Journal of Species Lists and Distribution* **10**: 407–408.
- Santana, D.J., Mângia, S., Silveira-Filho, R.R., Barros, L.C.S., Andrade, I., Napoli, M.F., Juncá, F., Garda, A.A. (2015): Anurans from the middle Jaguaribe River region, Ceará state, Northeastern Brazil. *Biota Neotropica* **15**: 1–8.
- Silva, L.A., Hoffmann, M.C., Santana, D.J. (2014): New record of *Corythomantis greeningi* Boulenger, 1896 (Amphibia, Hylidae) in the Cerrado biome, State of Tocantins, Central Brazil. *Herpetology Notes* **7**: 717–720.
- Silvano, D.L., Segalla, M.V. (2005): Conservation of Brazilian amphibians. *Conservation Biology* **19**: 6533–6538.
- Sueur, J., Aubin, T., Simonis, C. (2008): Seewave: a free modular tool for sound analysis and synthesis. *Bioacoustics* **18**: 213–226.
- Toledo, L.F. (2010): Description of a new species of *Pseudopaludicola* Miranda-Ribeiro, 1926 from the State of São Paulo, Southeastern Brazil (Anura, Leiuperidae). *Zootaxa* **2681**: 47–56.
- Toledo, L.F., Siqueira, S., Duarte, T.C., Veiga-Menoncello, A.C.P., Recco-Pimentel, S.M., Haddad, C.F.B. (2010): Description of a new species of *Pseudopaludicola* Miranda-Ribeiro, 1926 from the State of São Paulo, Southeastern Brazil (Anura, Leiuperidae). *Zootaxa* **2496**: 38–48.
- Vieira, W.L.S., Arzabe, C., Vieira, K.S. (2006): Amphibia, Leptodactylidae, *Leptodactylus caatingae*: geographical distribution extension. *Check List* **2**: 55–56.