

## Predation of *Mussurana bicolor* (Serpentes, Dipsadidae) upon *Leptodactylus podicipinus* (Anura, Leptodactylidae) in the Pantanal, Brazil

Diego José Santana<sup>1,\*</sup>, Adriana Maria Espinoza Fernando<sup>1</sup>, Aline da Silva Giroux<sup>1</sup>, Camyle Arye Maruyama<sup>1</sup>, Felipe Luis Gomes Borges<sup>1</sup>, and Isabel Melo<sup>1</sup>

Studies on snake trophic ecology are important to understand several of their evolutionary traits (Kardong et al., 1996; Sherratt et al., 2018). However, reports on predation events are dependent on fortuitous observations in the wild due their secretive habits, and dependence of such observations limits the understanding of snake trophic ecology (Bernarde, 2007). Snakes of the tribe Pseudoboini have been the focus of natural history studies (e.g., Alencar et al., 2013; Gaiarsa et al., 2013), which have shown a general tendency to prey on lizards and snakes. Therefore, most of the *Mussurana* species present ophiophagous behavior, but data on the diet of *Mussurana bicolor* (Peracca, 1904) are scarce, and the few records regarding its trophic ecology suggest it as one of the few pseudoboinine that feeds on amphibians (Gaiarsa et al., 2013). *Mussurana bicolor* is distributed in Southwestern Brazil, Paraguay and North of Argentina, and it occurs in open areas of the Pantanal and Chaco regions (Gaiarsa et al., 2013). *Leptodactylus podicipinus* (Cope, 1862) is a small and widely distributed leptodactylid frog, occupying open formations of Paraguay, Argentina, Bolivia, northwestern Uruguay, and central Brazil (Frost, 2018). It has been recorded as prey for some vertebrate species (e.g., Briso et al., 2014; Oda et al., 2014), showing its important role in the trophic chain where it occurs. In this work, we observed an attempt of predation of *Mussurana bicolor* upon the frog *Leptodactylus podicipinus*.

During field work on Base de Estudos do Pantanal (-19.576578°S, -57.018904°W), municipality of Corumbá, state of Mato Grosso do Sul, Brazil, we observed an individual of *Mussurana bicolor* (IBSP 90288, snout-vent length [SVL] 425.00 mm) in a predation attempt upon a female of *Leptodactylus podicipinus* (ZUFMS-AMP 11333, SVL 37.90 mm). This attempt of predation was observed at 21:00h in a permanent swamp. When we found this predatory event, the snake was biting the frog in the posterior members while constricting it (Fig. 1).

*Mussurana* snakes are well known by their ophiophagous predatory habits, feeding on different terrestrial, semi-aquatic and fossorial snakes (Alencar et al., 2013).



**Figure 1.** Predation attempt of *Mussurana bicolor* (IBSP 90288) upon *Leptodactylus podicipinus* (ZUFMS-AMP 11333) in the Pantanal, Base de Estudos do Pantanal, municipality of Corumbá, state of Mato Grosso do Sul, Brazil.

<sup>1</sup> Universidade Federal de Mato Grosso do Sul, Instituto de Biociências, Bairro Universitário, 79070-900, Campo Grande, Mato Grosso do Sul, Brazil.

\* Corresponding author. E-mail: jose.santana@ufms.br

Despite the few records on *M. bicolor*'s diet in the same studied area, there is a record of it preying upon the water snake *Helicops leopardinus* (Fiorillo *et al.*, 2013), reinforcing the ophiofagous habit of *Mussurana* species, even in the Pantanal wetland. *M. bicolor* is one of the few species of the genus feeding on amphibians (Gaiarsa *et al.*, 2013), showing that this species has a wider range of prey than its congeners.

The present report is the first record of *L. podicipinus* as prey of *M. bicolor*, and the fourth of an anuran as its prey (Strüßmann, 1992; Yanosky *et al.*, 1996; Gaiarsa *et al.*, 2013). Given the high abundance of *L. podicipinus* in the Pantanal, its common occurrence throughout the year (Prado *et al.*, 2005), this observed predator-prey interaction between *M. bicolor* and *L. podicipinus* might suggest that it is frequent in the Pantanal, where both species co-occur.

**Acknowledgments.** This report is result of a research conducted during the 21th Pantanal Field Course (EcoPan2018) of the Universidade Federal of Mato Grosso do Sul. We thank Liliana Piatti for her pre peer-review. We also thank Felipe Grazziotin for a previously review. AMEF and FLGB thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for scholarships. ASG, CM and IM thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for scholarships. DJS thanks CNPq for his research fellowship 430 (311492/2017-7).

## References

- Alencar, L.R., Gaiarsa, M.P., Martins, M. (2013): The evolution of diet and microhabitat use in Pseudoboine snakes. *South American Journal of Herpetology* **8** (1): 60–66.
- Bernarde, P.S. (2012): Anfíbios e répteis: introdução ao estudo da herpetofauna brasileira. Curitiba, Brazil, Anolis Books.
- Briso, A.L.F., da Graça, R.J., de Oliveira, M.R.F., Oda, F.H. (2014): Predation on pointbelly frog *Leptodactylus podicipinus* (Anura: Leptodactylidae) by the rufescent tiger-heron *Trigrisoma lineatum* (Pelecaniformes: Ardeidae). *Herpetology Notes* **7**: 731–732.
- Frost, D.R. (2018): Amphibian species of the world: an online reference. Version 6.0. Available at: <http://research.amnh.org/herpetology/amphibia/index.html>. Accessed on 14 September 2018.
- Gaiarsa, M.P., Alencar, L.R.V., Martins, M. (2013): Natural history of Pseudoboine snakes. *Papéis Avulsos de Zoologia* **53** (19): 261–283.
- Kardong, K.V., Kiene, T.L., Bels, V. (1996): Evolution of trophic systems in squamates. *Netherlands Journal of Zoology* **47** (4): 411–427.
- Oda, F.H., Oda, T.M., Felismino, M.F. (2008): *Leptodactylus podicipinus* (Pointedbelly Frog)-Predation. *Herpetological Review* **39**: 337–338.
- Prado, C.P.A., Uetanabaro, M., Haddad, C.F. (2005): Breeding activity patterns, reproductive modes, and habitat use by anurans (Amphibia) in a seasonal environment in the Pantanal, Brazil. *Amphibia-Reptilia* **26** (2): 211–221.
- Sherratt, E., Rasmussen, A.R., Sanders, K.L. (2018): Trophic specialization drives morphological evolution in sea snakes. *Royal Society open science* **5** (3): 172141.
- Strüßmann, C. (1992): Serpentes do Pantanal de Poconé, Mato Grosso: Composição faunística, história natural e ecologia comparada. Unpublished Dissertation, Universidade Estadual de Campinas, Campinas, Brazil.
- Yanosky, A., Dixon, J., Mercolli, C. (1996): Ecology of a snake community of El Bagual Reserve, Argentina. *Herpetological Natural History* **4**: 97-110.